# INSTITUTIONS

OF

# MEDICINE.

PART I.

### PHYSIOLOGY.

# For the use of the Students in the University of Edinburgh.

#### È D I N B U R G H:

M, DCC, LXXII.

### ADVERTISEMENT.

The following sheets are neither correct enough for the public eye, nor full enough for the most part of readers; but the apology is, that they are intended only for those who are to hear them fully explained.

# INSTITUTIONS

Q.F

# MEDICINE

I. Medicine is the art of preventing and of curing difeafes.

II. Before confidering the application of this art to particular difeases, certain general doctrines are neceffary to be premised, which are called THE INSTITUTIONS OF MEDICINE.<sup>\*</sup>

A 3

# ( 6 )

III. The Inftitutions of Medicine: are divided into three parts.

The first treats of life and health.

The fecond delivers the general doctrine of difeafes.

The third delivers the general doctrine concerning the means of

### preventing and curing difeafes.

INSTI-

# INSTITUTIONS of MEDICINE.

# PART I. PHYSIOLOGY..

IV. THE doctrine which explains the conditions of the body and of the mind neceffary to life and health, is called PHYSIOLO-GY, or the Doctrine of the Animal OEconomy.

V. The functions of the animal oeconomy are many and various, and fo complicated with each other, that it is difficult to find the most proper order in which they may be delivered. That, however, feems the best which confiders them as nearly

as may be according to the feries of causes and effects.

VI. Upon this plan, we fhall treat.I. Of the folid matter, of which a great part of every organ of the body confifts.

2. Of the nervous fystem, in which the motions of the body for the most part begin, and upon which the motions produced in it chiefly

depend.

3. Of the motion and circulation of the blood, and of the feveral organs and actions employed in fupporting it.

4. Of the functions employed in fupporting and repairing the feveral folid and fluid matters of the body; and, on this occasion, of the nature of the feveral fluids themfelves.

5. Of

5. Of the organs employed in receiving and modifying the impreffions of external bodies neceffary to fenfation; and of their feveral func-tions.

6. Of the motions of the whole body, or of its feveral parts, which depend on the action of museles, and not before explained.

7. Of the functions peculiar to the fexes, and of generation.

PHY-

### SECT. I.

OF THE SIMPLE SOLIDS.

VII. The folid parts of the body feem to be of two kinds; one whole properties are the fame in the dead as in the living, and the fame in the animate as in many inanimate bodies; the other whole properties appear only in living bodies. In the last, a peculiar organization, or addition, is fupposed to take place; in opposition to which, the first are called the SIMPLE SO-LIDS. Of these only we shall treat here, and of the others, which may be called VITAL SOLIDS, being the fundamental part of the nervous fystem,

fystem, we shall treat under that title in the following section.

VIII. The fimple folids are fuitcd to the purpofes of the animal oeconomy by a certain force of cohefion, joined with a certain degree of flexibility and elafticity.

IX. These properties of the fimple folids, in different parts of the

body, in different bodies, and on different occasions in the fame body, are neceffarily in different degrees; and this feems to depend upon the difference of the mixture, aggregation, or organization of the folid.

X. The matter of the fimple folid every where, except in the bones, appears to be an homogeneous ag<sup>2</sup> gregate;

gregate; and there is no proper evidence of its being formed of certain parts naturally difcrete and incoherent, which are cemented by others of a different nature.

XI. Of the fimple folid confidered as an homogeneous aggregate, the integrant parts are a mixt, which feems to be nearly of the fame kind in all the different parts of the hu-

man body, and perhaps in moft of the parts of every animal: So far as we yet know, the variety of it is very inconfiderable.

XII. This, which may be called the ANIMAL MIXT, is found, by chemical experiment, to be confiderably different from every kind of vegetable or foffil matter; but the fame experiments hardly teach us any

anything exact or useful with respect to the constituent parts of this mixt.

XIII. The only particular relative to this, which we exactly know, is, that the animal mixt is formed of water, and of fome other matter concreting with it; that, on different occafions, the ftate of it is varied by the proportion which the water bears to the other concreting matter; and that, efpecially by a different proportion in this refpect, the fimple folid differs in its force of cohefion, flexibility, and elafticity (VIII).

XIV. The proportion of water to the other matter in the animal mixt of different perfons feems to depend, in the first place, upon the B nature

nature of the original flamina in each; as the different flate of fimple folids, which appears early to diftinguifh fex and temperament, continues refpectively the fame thro' the whole of life, even though the different perfons are under the fame external circumflances.

XV. But, in every particular perfon, that proportion is conftantly changed by the progrefs of life; and this happens more or lefs as other caufes concur.

XVI. The caufes that can affect the mixture of the fimple folid, are either the flate of the nutritious fluid conveyed by the ordinary channels, or fome matters from without infinuated into the folid.

XVII.

XVII. The flate of the nutritious fluid may be varied by the quantity and quality of the aliment taken in, by the powers of concoction and affimilation, by the circumflances of application and concretion, or by certain practernatural matters cartied along with it.

XVIII. The external matter that may be infinuated into the fimple folid is various, but for the moft part is only aqueous moifture in greater or lefs quantity.

XIX. That these feveral causes may affect the proportion of water in the fimple folid, and thereby give a different state of it, is sufficiently obvious: That the same causes may also affect the other concreting matter, we can, in general, perceive **B** 2 to

to be poffible; but in what manner, or upon what occasions they do so, is not easily differed.

XX. The properties of the fimple folid (VIII.) may be also varied by its state of aggregation; and this again may be varied, 1. By the temperature of the atmosphere to which the body is long exposed. 2. By the preliure, external or internal, which is applied to the folid. 3. By the degree of extension of the folid beyond its natural state, which, in every living body, is given more or lefs to every part of the foft or flexible folids; and, lastly, by the motion or reft to which the folid is accuftomed.

XXI.

XXI. The properties (VIII.) of the folid parts are alfo varied by the flate of their organization. This every where depends upon an arrangement of fibres, the flate of cellular texture, or upon a texture of veffels; and therefore, to explain the different flates of organization, it will be enough to mention the caufes of the differences which occur in thefe fundamental parts.

XXII: Fibres may differ in fize by the feveral caufes (XIV.--XXI.) affecting the mixture aud aggregation of the matter of which they are formed, and by thefe caufes alone; but how far the organization of any part depends upon an arrangement of fibres, we cannot diffinctly perceive; and, if it does, we cannot perceive that the flate of fuch parts differs o-B 3 therwife

therwife than by the ftate of the cellular texture every where interpofed between the fuppofed fibres.

XXIII. The ftate of cellular texture is the most important circumstance in all organized parts ;. and it may be varied by many different causes. 1. The texture may be more denfe, and thereby firmer, as it has been more. preffed by the actions of life or external force; by which means especially it is changed in the progress of life. 2. The cellular texture may be increased in bulk, and rendered firmer by a new growth taking place in it, as frequently happens in membranes which are flowly and gradually ftretched out. 3. The fame texture may become weaker by fome part of it being eroded

eroded by acrid matters generated in the body, or externally applied. 4. It is analogous to this, that, when any part is fustained by feveral. layers of cellular texture or membranes, such support is weakened. by one or more of these layers being cut through; and the fame weaknefs is induced when any external compression, which, for fome time, had been applied, is taken away. 5. The flate of the cellular texture is varied by the matter contained in its cells; which is fometimes a matter concreting into a folid mass, and sometimes a preternatural quantity of an aqueous inelastic fluid. The bones formed in the first manner may again become foft by the hardened. matter's being diffolved and reabforbed. 6. When the mobility.

ty of parts on one another depends upon the extent of cellular texture connecting them, that mobility is diminished or destroyed by a great part of the cellular texture being eroded or cut away, and the remaining parts being united together; fo that the parts are now connected by a shorter portion of cellular texture than before. 7. Parts naturally separate may lose their mobility by being joined together by a cellular texture formed between them, as happens when any two furfaces are for some time kept closely applied to each other:

XXIV. In so far as a solid part is formed by a texture of vessels, its properties (VIII.) may be varied by the different states of these vessolutions fels;

fels; which, 1. may be more or lefs full of fluids. 2. They may be changed into a folid mafs, by the fluid, contained and flagnating in them, concreting into a folid. 3. They may be changed into a folid, if the fluids that fhould pafs through them are intercepted, and the cavity is filled with a cellular texture ; or, 4. They may be changed into a folid, if, by collapfe or preffure, the

fides of the veficls are applied to each other, and concrete together.

XXV. The pathology of the fimple folids cannot be properly feparated from their phyfiology; and therefore, many different flates of thefe folids, though fuch as are always morbid, are mentioned above. We think it alfo proper to fubjoin here

here a fhort view of the whole of that pathology.

XXVI. The difeafes of the fimple folids are,

I. Those of the naturally fost parts.

J. Debility with flexibility.
 Debile tenerum, gracile, Gaubi
 Pathol. 161. 1.
 Debile tabidum Gaub. ibid. 161. 2.

 Debility with fragility. Debile fiffile, Gaub. 161. 3.
 Laxity. Debile laxum, flaccidum, Gaub. 160. 1.
 Flaccidity. Debile iners, Gaub. 160. 2.
 Rigidity diminifhing flexibility.

Rigi-

# PHYSIOLOGY. 23.

Rigidum tenax, Gaub. 165. 1. 6. Rigidity deftroying flexibility. Rigidum durum, Gaub. 165. 2.

- II. Those of the naturally hard parts.
  - Flexibility.
    - Debile flexile, Gaub. 160. 3.
  - 2. Fragility.

Fragile fpongiofum, Gaub. 161. 4. Fragile vitreum, Gaub. 165. 3. Of all these morbid affections, we suppose the remote and proximate causes may be understood from what is delivered above, (XIV.— XXIV.)

#### SECT.

#### SECT. $\Pi$ .

#### OF THE NERVOUS SYSTEM.

XXVII. The nervous fyftem, as the organ of fenfe and motion, is connected with fo many functions of the animal oeconomy, that the fludy of it must be of the utmost importance, and a fundamental part of the fludy of the whole oeconomy.

# A general View of the Nervous System.

XXVIII. The nervous fyftem confifts of the medullary fubftance of

of the brain, cerebellum, medulla oblongata, and fpinalis; and of the fame fubftance continued into the nerves, by which it is diffributed to many different parts of the body.

XXIX. The whole of this fyftem feems to be properly diffinguished into these four parts.

1. The medullary fubstance con-

tained in the cranium and vertebral cavity, the whole of which feems to confift of diffinct fibres, but without the feveral fibres being feparated from each other by any evident enveloping membranes.

> N. When we speak of functions, which are or may be in common to every part of this portion of the nervous system, we shall speak of the whole under the title of the BRAIN; C

BRAIN; but, when it is necessary to diffinguish the particular parts, we shall take care to avoid ambiguity.

2. Connected with one part or other of No. 1. are, the NERVES, in which the fame medullary fubstance is continued, but here more evidently divided into fibres, each of which is separated from the others by an enveloping membrane derived from the pia mater. 3. Parts of the extremities of certain nerves (2.) in which the medullary substance is divested of the enveloping membranes from the pia mater, and fo fituated as to be exposed to the action of certain external bodies, and perhaps fo framed as to be affected by the action of certain bodies only: These we name the SENTIENT EXTREMI-TIES of the nerves.

4. Certain extremities of the nerves (2.) fo framed as to be capable of a peculiar contractility, and, in confequence of their fituation and attachments, to be, by their contraction, capable of moving most of the folid and fluid parts of the body. These we name the MOVING EXTRE-MITIES of the nerves: They are commonly named MOVING or MUSCULAR FIBRES.

N. That the mufcular fibres are a continuation of the medullary fubftance of the brain and nerves, has not been fhewn by the anatomifts, nor univerfally admitted by the phyfiologifts; but we now fuppofe it, and hope afterwards to render it fufficiently probable.

Are the ganglions of the nerves to be confidered as a part of the ner-C 2 vous



# 28 PHYSIOLOGY. vous fystem diffinguished by a peculiar function?

XXX. These feveral parts of the nervous fystem are every where the fame continuous medullary fubstance which we suppose to be the vital folid of animals, so constituted in living animals, and in living fystems only, as to admit of motions being readily propagated from any one part to e-

very other part of the nervous fyftem, fo long as the continuity and natural living flate of the medullary fubflance remains.

N. It is obferved, that the comprelion of any part of the medullary fubflance prevents the communication of motion between the parts that lie on different fides of the part comprefied; and it is probable, there are other caufes befides comprelion, which may also affect the medullary fubflance.

flance, fo as to interrupt in it the communication of motion; but they are not diflinctly known. In the mean time, we use the expression, of a nerve, or other portion of the nervous fystem being *free*, to denote its being free, not only from compression, but from every other suppofed cause interrupting the communication of motion.

The condition fitting the medullary fubfance for having motion propagated in it, we fuppole to be the prefence of a certain fluid; which we therefore name the *nervous fluid*, wit'tout meaning however at prefent to determine any thing with regard to itsfource, nature, or manner of acting.

XXXI. In the living man, there is an immaterial thinking fubftance or MIND conftantly prefent; and every phaenomenon of thinking is to be confidered as an affection or faculty of the mind alone. But this immaterial and thinking part of man is fo connected with the material and C 3 corporeal.

corporeal part of him, and particularly with the nervous fyftem, that motions excited in this give occafion to thought; and thought, however occafioned, gives occafion to new motions in the nervous fyftem. This mutual communication or influence we affume with confidence as a fact: But the mode of it we do not underftand, nor pretend to explain; and therefore are not engaged to obviate the difficulties that attend any of the fuppofitions which have been made concerning it.

XXXII. The phaenomena of the nervous fystem occur commonly in the following order. The impulse of external bodies acts upon the fentient extremities of the nerves; and this gives occasion to preception or thought, which,

which, as first arifing in the mind, we term SENSATION. This fensation, according to its various modification, gives occasion to VOLITI-ON, or the willing of certain ends to be obtained by the motion of certain parts of the body; and this volition gives occasion to the contraction of muscular fibres, by which the motion of the part required is produced.

N. This is an example of the most ordinary cafe; but we do not mean to fay. It is the only cafe of communication between the different parts of the nervous fystem:

XXXIII. As the impulse of bodies on the fentient extremities of a nerve does not occasion any fensation, unless the nerve between the fentient extremity and the



the brain be free (XXIX. 3.); and as, in like manner, volition does not produce any contraction of mufcles, unlefs the nerve between the brain and muscle be also free; we conclude, from both these facts, that fenfation and volition, fo far as they are connected with corporeal motions, arc functions of the brain alone; and : we presume, that sensation arises only in confequence of external. impulse, producing motion in the sentient extremities of the nerves, and of that motion's being thence propagated along the nerves to the brain; and, in like manner, that the will operating in the brain only, by a motion begun there, and propagated along the nerves, produces the contraction of muscles, XXXIV.

XXXIV. From what is now faid, we perceive more diffinctly the different functions of the feveral parts of the nervous system, as distinguished in (XXIX.) I. The fentient extremities (XXIX. 3.-) feem to be particularly fitted to receive the impreffions of external bodies, and according to the difference of these impreffions, and of the condition of the sentient extremity itself, to propagate along the nerves motions. of a determined kind, which, communicated to the brain, give occasion to fenfation. 2. The brain (XXIX. 1.) feems to be a part fitted for, and susceptible of those motions with which fenfation, and the whole consequent operations of thought are connected, and thereby is fitted to form a communication between the motions

motions excited in the fentient, and those in consequence arising in the moving extremities of the nerves, which are often remote and diftant from each other. 3. The moving extremities (XXIX. 4.) are fo framed as to be capable of contraction, and of having this contraction excited by motion, propagated from the brain, and communicated to the contractile fibre. 4. The nerves, more strictly fo called (XXIX. 2.), are to be confidered as a collection of medullary fibres, each inveloped in its proper membrane, and thereby fo feparated from every other, as hardly to admit of any communication of motion from any one to the others, and to admitonly of motion along the continuous medullary substance of the same fibre,

PHYSIOLOGY. 35 fibre, from its origin to the extremitics, or contrariwife.

XXXV. From this view of the parts of the nervous system, of their several functions and communication with each other, it appears, that the beginning of motion in the animal æconomy is generally connected with fensation; and that the ultimate cffects of fuch motion are chiefly actions depending immediately upon the contraction of moving fibres, between which and the fentient extremities, the communication is by means of the brain. Wherefore, in ftudying the nervous fystem, we judge it proper to consider, 1. Senfation, and with that the general function of the sentient extremities. 2. The action of the moving fibres. 3. The



3. The function of the brain. In confidering these three heads, the function of the nerves, more strictly fo called, will, of course, be sufficiently explained.

#### CHAP. I.

OF SENSATION.

XXXVI. Our fenfations may be confidered as of two kinds: 1. Thofe ariting from the impulfe or imprefiion of external bodies, which we therefore name SENSATIONS OF IM-PRESSION. 2. Thofe arifing from the mind's being confeious of its own action, and of the motions it excites; and thefe we name SENSATIONS OF CONSCIOUSNESS.

Sen-

Sensations of Impression.

XXXVII. The fenfations of imprefion are very various, but have been generally referred to five heads or kinds, commonly called the five fenfes; that is, those of fight, hearing, fmell, tafte, and touch.

XXXVIII. The four first of these are each of them properly confidered, as forming one genus of fensations : I. As the particular fensations comprehended under each head (XXXVII.), though very various, are, however, perceived to have fomewhat common to all of them. 2. As those of the same genus all arise from impressions made D upon

upon one part of the body only, and that of a peculiar organization. 3. As those of the fame genus all arife from the action of external bodies of one kind only, or of one and the fame quality, by means of which they act upon our organs.

XXXIX. No fuch characters concur in eftablishing one genus of the fenfations referred to the fifth head

of touch, which are various in all those respects (XXXVIII.); and physiologists feem to have referred, to this head of touch, every fensation that does not manifestly belong to the other four; and, among the rest, many of the sensations of conficious fees. It might perhaps be useful to distinguish into genera, the several sensations referred to touch; but

but it is not necessary to be done here.

From certain fenfations referred to touch, it appears, that not only the extremities (XXIX. 3.), but that every part of the nervous fyftem (XXVIII.) is fentient with respect to certain impressions.

Sensations. of Consciousness.

XL. The fenfations of confcioufnefs may be referred to the following heads: 1. Thofe of apperception, by which we are in general confcious of thinking, of perceiving, judging, and willing, and thereby of our existence and identity. 2. The fenfations arising from D 2 the



the particular state of thinking, as perception, memory, and judgement, are more or less clear, ready, or exact. 3. The fensations arising from the particular flate of volition, and its various modes. 4. The fenfations arising from the general state of action, as vigorous or weak, eafy or difficult. 5. The fensations arifing from particular actions, or a confciousness of the actions excited, and of the motion of the different parts of the body. 6. The fenfations arising from the diminution or absence of impressions.

Under each of these heads, a great number of particular sensations are comprehended, but not necessary to be farther specified here.

Laws,

Laws, or general Circumstances of Senfation.

XLI. Of the four first genera (XXXVII.), the fentations arifing give no indication of the nature of the bodies acting on our organs, or of the mode of their action; and, when we otherwife learn these cir-

cumftances, we can perceive no neceffary connection between them and the fenfations which they produce. But, from certain fenfations of touch and confcioufnefs, we acquire the notions of folid figure, of motion, impulfe, impenetrability, and the communication of motion, and confider the fenfations as exactly correspondent to the circumftan-D.3 ces



ces of external bodies. At the fame time, as we know of no other action of bodies on each other, but that of impulfe; and as, in the cafe of the fenfations of the four firft genera, we learn, that an impulfe takes place, we have comprehended the whole under the title of fenfations of imprefiion, and confider all of them as perceptions of impulfe.

XLII. To produce any fenfation of impreffion, a certain force of impreffion is neceffary; and, from a leffer force, no fenfation arifes. The degree of force is alfo limited on the other hand; as, in a high degree, it deftroys the organ; and, in degrees approaching to this, a general fenfation of pain, rather than

than the sensation of any particular object, is produced.

XLIII. Within these limits, however, our sensations are not exactly correspondent to the force of impression, nor do they make any exact estimate of that force. For the most part, sensation is relative to the change that is produced in the nervous system, and a sensation

proves firong or weak, only as it is fironger or weaker than that which had immediately preceded it, or than that degree of force to which the nerves had been immediately before accuflomed. For this reafon too the limits (XLII.) are very variable.

# (XLIV,)

XLIV. Different fenfations do not always imply a different kind of action in the bodies producing them; for fometimes different fenfations arife merely from a different degree of force in the fame kind of action, as is manifest in the cafe of heat and cold.

XLV. To fendation from impreffion, a certain duration of impreffi-

on is necessary.

XLVI. The mind's refting for fome time upon one fenfation is called ATTENTION. This, like the duration (XLV.) is neceffary to give an imprefiion its full effect.

XLVII,

XLVII. The mind feems to be determined to attention by the force of impreffion; by the pleafure or pain arifing from it; by the degree of emotion or paffion produced by thefe; and, laftly, by the emotion's being more or lefs related to the perfon feeling.

XLVIII. If the force and duration of impreffion, and the attention of mind, are all in the due degree, the fenfation often remains for fome time after the impreffion or action of the external body has ceafed.

XLIX. The mind admits of, or or can attend to, one fenfation only at one time.



L. Though the mind admits but of one fenfation at one time, feveral imprefions may act at the fame time, if they be fuch as can unite in producing a fingle fenfation; and fuch is the cafe of many of the impreffions which produce the particular fenfations of the fame genus, as in those efpecially of colour, found, fmell, and tafte.

LI. In each of these genera, many impressions which separately produce particular species can unite in producing a single sensation which is always a neutral, or one different from either of the separate sensations.

LII. This union of imprefions may take place, either when the impreffions are exactly fynchronous, or

or when the one fucceeds the other before the fenfation of the firft (XLVIII.) has ceafed.

LIII. Though the motion excited in the fentient extremitics by imprefion remains fome time, as in (XLVIII.), it must be fupposed to become continually weaker, till at length it ceases altogether, and with it the fensation.

LIV. The fame impreffion foon repeated, does not produce the fame ftrength of fenfation as before. Hence, all new impreffions are, *caeteris paribus*, ftrongeft; and moderate impreffions frequently repeated produce no fenfation, unlefs their force is confiderably increafed.



LV.

LV. Actions which at first produced a fensation of confcioufness, as accompanied with volition, come, by repetition, to be performed without any fensation; or they produce it only when they are performed with uneafines, pain, or unufual force.

LVI. Impressions being given, their effects in producing fensation

are different in different perfons, and in the fame perfon at different times. This muft arife from fome difference in the ftate of the bodies acted upon, which may perhaps be referred to the following heads: 1. The flate of the common teguments, or other parts interpofed between the impreffing body and the medullary fubftance of the fentient extremity. 2. The

The different state of the medullary fubstance of the sentient extremities. as given to it in the original stamina. 3. The different state of tension in the medullary fubstance of the fentient extremities, as given to it by the state of the blood-vessels constantly connected with it. 4. The ftate of the fame medullary fubstance, as affected by heat or cold. 5. The ftate of it, as produced by former impressions (XLIII. LIV.) 6. The state of the nerves along which the motion is propagated. 7. The state of the brain or fensorium. 8. The state of attention (XLVI. and XLVII.)

LVII. Different parts of the body are fenfible, and fenfible only by means of nerves prefent in them; but anatomy does not always de-E termine

termine certainly with regard to the prefence or abfence of nerves; and, therefore, the fenfibility of feveral parts can be determined by experiment only; which, however, is alfo fallacious.

LVIII. Particular senfations arife from impressions on certain parts only: 1. Because the sentient extremities in these parts are so fituated as to be exposed to the action of certain external bodies only. 2. Becaufe the fentient extremities are connected with an organ that increases the force of the external agent, or modifies its action in the manner neceffary to a determined impression. 3. Because the fibres of certain sentient. extremities are, by their fize or tension, fitted to be acted upon by cer-

certain external bodies only. 4. Becaufe certain fentient extremities are fo conftantly preferved in a certain ftate, as to render them more fenfible to a change.

These circumstances deretmine the mode of impulse, but do not account for the sensation arising from it.

LIX. Different sensations are accompanied with different judgements

concerning the bodies making impreffion, and the part of the human body upon which it is made. Some fenfations are referred to bodies at a diftance; others, to external bodies in contact; and others to the feeling body itfelf.

LX. When fenfations are referred to our own bodies, it is in three fe-E 2 veral

1

veral ways: 1. They are most commonly referred to the part on which immediately the impression is made, and this, with regard to the external parts, very accurately; but, with regard to the internal, much less fo: And commonly the fenfations arising from internal parts, are referred to the incumbent external part, with some obscure distinction between superficial and more deep. 2. Senfations are sometimes referred, not to the part upon which the impression is immediately made, but to a diftant more fenfible part, to which a motion is propagated from the part impreffed. 3. As fenfations ufually arife form impreffions made upon the extremities of the nerves, and are referred to these, so impressions made on the nerves in their course, are fome-

fometimes referred to the extremities from whence they were in use to arife.

LXI. The fenfations of confcioufnefs (XL. 1. 2.) are referred to the *encephalon*. So are those of XL. 3. if they are moderate; but, if more vehement, they are often referred to those parts in which their effects are exerted, as the heart and organs of respiration. The sensations (XL. 4. and 5.) are feldom, with accuracy, referred to particular parts, but indistinctly to a whole member. We are not confcious of the action of particular muscles, except when their contraction is spafmodic.

LXII. We are difpofed to combine our fenfations, as united in one E 3 ob-



54 PHYSIOLOGY. object; and thus form what are called COMPLEX IDEAS.

LXIII. We compare our feveral fenfations, and from thence acquire new fenfations of RELATION.

LXIV. When fenfations formerly received are again renewed by the fame objects, it is, for the moft part, with a confcioufnefs of their having been formerly received; and this faculty we call REMINISCENCE.

LXV. Perceptions formerly received can be renewed without the prefence or action of the object which formerly gave occasion to them; and if this is attended with the confciousness of a difference between the vividity of the two perceptions, and

and particularly of the absence of the original objects, such a renewed perception is called an IDEA; and the faculty by which this renewal is made is called MEMORY.

LXVI. Perceptions formerly received, can, without the prefence of the original object, be renewed alfo in fuch a manner, that the mind does not perceive any difference between the original and the renewed perception; and therefore, fuch renewal is always with the perfuafion of the prefence of the object. The faculty by which fuch renewal is made, we call IMAGINATION, more ftrictly.

LXVII. Reminiscence depends upon the force or frequent repetition of the former sensation.

LXVIII,

## 56 PHYSÍOL OGY.

LXVIII. Memory depends upon an affociation of perceptions, which is formed by their being frequently repeated immediately after each other; by their being parts of the fame complex idea; and, by their having relations marked. Memory is generally faithful to fuch affociations; but it is more or lefs fo in different perfons, according to the number and importance of the relations marked,

according to the frequency of the repetition of the fenfations, and the marking of their relations; and according to the different states of the brain, very little known.

LXIX. Imagination feems always to depend upon internal causes, that is, upon causes acting in the brain.

LXX.

LXX. Memory and imagination renew diffinctly the ideas of feeing and hearing only. All others are renewed imperfectly, or not at all; but all others may be affociated LXVIII.) with the sensations or deas of feeing and hearing, fo that hese become signs of the others. The memory, in renewing these ligns, fo far renews the idea belonging to them, as to renew their feveal affociations and relations; to renew, in some degree, the pleasure or pain which formerly attended the enfations them felves, and particulary to renew the emotions of mind, motions of the body, which the enfations formerly produced.

LXXI.



LXXI. Moft of our fenfations, perhaps all of them, are either pleafant or painful.

LXXII. The words pleasant and painful are commonly generic terms, each of them comprehending a great many species, which seem to require being afforted under feveral different genera. Thus, in the firk place, our fensations may be divided into these we defire, and those we are averse to. Of these we defire, we may diffinguish those arifing from qualities we refer to othe bodies, from those we refer entirely toourown. Thefirst may be name more ftrictly the AGREEABLE, the last the PLEASANT. In like manner, of the sensations we are aver tG

to, we may diffinguish the DISA-GREEABLE and the PAINFUL. But, farther, the last must be distinguished from that sense of aversion which accompanies certain sensations of confciousnels, as the sense of debility, lassitude, difficulty, &c. and, particularly, from that referred obscurely to internal parts, which we name ANXIETY. These fenfations may be called the UNEASY; and every one diffinguishes this kind from that of the PAINFUL, more strictly fo called. These last feem to be always fenfations of impreffion, referred pretty accurately to a particular part. There is thus a foundation for e-Rablishing different genera of the fenfations we defire, and of those we are averse to; as also, for greater.

pre- .



precifion in the employment of terms: But the fixing the limits of these genera, and afforting the several species, may be still difficult; so that we cannot be certain of applying the terms every where with strict propriety.

LXXIII. The enumeration of the agreeable or difagreeable, and even of the pleafant fenfations, would not be of much ufe here; and the enumeration of the uneafy and painful, though much more interefling, belongs to the pathology. However, we think it proper to deliver here the few following propositions.

LXXIV. Senfation and action, within certain limits, are always defired; and the want of fenfation, or imperfect



imperfect and indiffinct fenfations, are always uneafy. In action of every kind, the fenfations of debility and difficulty are also uneafy.

LXXV. In fenfations of impreffion, their being pleafant or painful often depends on the degree of force in the imprefion, allowance being made for the fenfibility of the fyftem.

LXXVI. As imprefiors, by being repeated, produce weak fenfations (LIV.), imprefions, at firft painfull, may, by repetition, be changed into pleafant, and the pleafant into infipid and uncafy. Hence arifes, with regard to moderate impreffions, the pleafure of novelty, the defire of variety, and the defire of in-F creafing

# 62 PHYSIOLOGY. creasing the force of pleasant impreflions.

LXXVII. There is a condition of imprefiions rendering them objects of defire or averfion, that cannot, with certainty, be referred to their force. This condition we call the quality of impressions.

LXXVIII. Imprefions are often redered objects of defire or averfion, by combination, fucceffion, and relation.

LXXIX. No fenfations arife originally in the mind, without a previous change in the flate of the body.

LXXX

LXXX. Certain imprefiions, and certain flates of the body, like to thofe which produce the fenfations of confcioufnefs, may both of them act upon the nervous fyftem, without producing any fenfation.

# SECT. II.

CHAP. II.

#### OF THE ACTION OF MOVING FIBRES.

LXXXI. The moving fibres (XXIX. 4.), fo far as yet known, are of one kind only, and the fame every where, as in the most commonly known muscles. Hence, the terms moving and muscular fibres are of the fame import.

F 2

LXXXII.

LXXXII. A mulcular fibre is fuppofed to have a peculiar organization, different both from that of the imple folid fibre, and from that of the medullary fibres in every other part of the nervous fystem; but wherein that peculiarity of organization confist, is not yet afcertained.

LXXXIII. A muscular fibre is

endued with a contractility, which is different from that of the fimple folids, or of any inanimate elaftics, efpecially in this, that the contraction of a mufcular fibre is excited by caufes which do not affect thefe others. For, the contraction of a mufcular fibre is excited by being extended; and a contraction is produced, whilft the ftretching power continues

continues to be applied. The fame contraction is alfo excited by various applications, whofe mode of action we do not perceive ; but we know them to be fuch as do not affect inanimate elaftics.

In refpect of these causes by which it may be excited, the contractility of muscular fibres hath been called IRRITABILITY.

LXXXIV. The force of contraction in mulcular fibres is often much greater than that of the caules exciting it.

LXXXV.The contactility ofmufcular fibres (LXXXIII.) appearsefpecially in living bodies, ceafeswith life, or foon after, and is  $pro-F_3$ F 3



bably never produced but with life. Hence it is, that, by fome writers, it is called the VITAL POWER, and the mufcular fibre endued with it, the LIVING SOLID. *Gaub. Patholog.* 169. 170.

LXXXVI. The contractility (LXXXIII.LXXXIV. & LXXXV.) hath been fuppofed to belong to mufcular fibres, independently of their connection with the other parts of the nervous fyftem; and, upon that fuppofition, it hath been called the *vis infita*. We fhall call it the INHERENT POWER; *Haller. Prim. lin.* 400.

LXXXVII. The contraction of inufcular fibres can be excited by applications made to other parts of the nervous

nervous fystem, as well as to the muscles themselves; and, as the effects of those applications made to the other parts of the nervous fystem can be prevented by ligatures made upon the nerves between the place of application and the muscle to be moved, it is concluded, that the constraction of muscular fibres can be excited by a power communicated to them by a motion propagated along the nerves. This power is called the NERVOUS POWER.

Sin.

LXXXVIII. The nervous power (LXXXVII.) is commonly determined to motion by the will. This we suppose to act in the brain only (XXXIII.), and to depend upon fienfation, and other modifications of thought; and this power, which is



to.

to be chiefly referred to the mind and acts in the brain only, we name the ANIMAL POWER.

LXXXIX. The facility with which the contraction of mulcular fibres can be excited, and the force with which it can be performed, are to be diftinguished. The first we name the MOBILITY, the last, the VIGOUR of mulcular fibres. Both have been confounded under the name of Irritability.

XC. The mobility and vigour of mulcular fibres (LXXXIX.) can both of them be increased or diminished by various means. Whatever can excite the contraction of mulcular fibres is called a STIMULUS; and, in

ingeneral; the means of exciting contraction are called STIMULANT POWERS. The means of diminifhing the mobility and vigour of mufcular fibres are called SEDATIVE POWERS.

XCI. The inherent power (LXXXVI.) is fuppoled to be more vigorous, moveable, and permanent in certain mufcular fibres than in

others.

XCII. The inherent power, or the contraction dependent upon it, can be excited by certain applications made either to the mufcles themfelves, or to the nerves connected with them; and, in either cafe, the effects of fuch application are fo exactly the fame, as to allow us to conclude, that the



the matter in the nerves, and in the muscular fibres, is of the same kind.

XCIII. The muscular fibres are fensible to various impressions, and are otherwise organs of the fenfations of consciousness (XL. 4. 5.) From this alfo, it is prefumed, that the mulcular fibres confift of the fame mater which is the fubject of fenk in other parts of the nervous fystem (XXXIX:)

XCIV. From (XCII. XCIII.) and other confiderations, we think it probable, that the muscular fibres an a continuation of the medullary fub stance of the brain and nerves, as alledged (XXIX.)

XCV. Though the muscular fibre consistof the same kind of matter # that in the nerves, the latter shew no con.

contractility, becaufe they have not the peculiar organization (LXXXII.) of the former.

XCVI. The nervous power (LXXXVII.), and the inherent (LXXXVI.) may fubfift for fome time without any connection of the nerves or mufcles with the brain; and they fubfift alfo in intire bodies for fome time after life has feemingly ceafed. Both powers, however, are feemingly of equal duration in thefe refpects; and neither power feems to fubfift long but in intire and living bodies.

XCVII. From what is faid (XCII.--XCVI.), it is probable, that the nervous and the inherent powers are fomewhat of the fame nature; and



and it is alfo probable, that, in intire and living bodies, both the nervous and inherent powers have a conflant dependence upon the animal (LXXXVIII.)

XCVIII. The contraction of mufcular fibres does not depend immediately on the motion of the blood, as it fubfifts, in many animals, after all motion of the blood has ceafed.

XCIX. The contraction of mulcular fibres does not depend on the inflation of vehicles, or other fuch analogous flructure, as the flortening of the fibres in contraction is often greater than can take place in fuch ftructure.

C. As the force of cohelion in the muscular fibres of living animals is much greater than in those of dead ones, it is probable from this, and other confiderations, that the caufe of muscular contraction is an increase only of that same power which gives the contractility of the fimple folids, and of other inanimate elastics; Haller. Prim. Lin. 407.408. If this is true, it will also explain why the force of cohefion in muscular fibres is greater than that of the medullary fibres in any other part of the nervous system, though both kinds of fibres, by (XCIV.) confift of the fame Lind of matter.

CI. In living and healthy animals, the mufcular fibres have a con-G ftant



conftant tendency to contract; and this tendency we call their TONE, or TONIC POWER.

CII. The tonic power of mufcular fibres neceffarily fuppofes their being conftantly in a ftate of extenfion beyond their natural or moft contracted ftate; and in this ftate they are conftantly kept by the action of antagonift mufcles, by the weight of the parts they fuftain, by fluids diftending the cavities they furround, and by their connection with fuch diftended cavities, particularly the blood-veffels.

CIII. As the differition of mulcular fibres, by (LXXXIII.), proves a ftimulus (XC.), we conclude, that the tonic power in them will, caeteris

teris paribus, be' in proportion to the degree of tenfion (CII.)

CIV. If the inherent power, as in (XCVII.), is in dependence upon the nervous and animal powers, and these may be increased or diminished by various means, the tonic, as a part of the inherent power, must, in some measure, be in proportion to the state of the nervous and ani-

mal powers.

CV. If the tonic power of any mulcular fibre depends more upon its flate of tenfion (CIII.) than upon the flate of the nervous and animal powers (CIV.) fuch fibres will be more affected by changes of the flate of tenfion, than by changes in  $G_2$  the

the flate of the nervous and animal powers; and on the contrary, &c.

CVI. The force of contraction, or the vigour of mulcular fibres, will be always as the force of ftimulus, and the vigour of the animal, nervous, and inherent powers taken together.

CVII. The mobility of muscular

fibres (LXXXIX.) feems often to be increafed by caufes which weaken their vigour; and, therefore, it is induced by the diminution of tenfion, and by caufes weakening the animal, nervous, or inherent powers.

CVIII.

CVIII. The ordinary contraction of mulcular fibres is disposed fpontaneously to alternate with a relaxation or extension of the same.

CIX. In the ftraight mufcles, and in the heart, the alternate contractions and relaxations readily take place; and that though a ftimulus is conftantly applied; but in mufcular fibres furrounding cavities, as in the alimentary canal, bladder of urine, &c. the alternate motions do not appear, unlefs a portion of the fibres is cut out, and feparated from the reft.

CX. From a difference in the ftate of a mufcle contracted by inherent power, while the member it fuftains is moved by external force,  $G_3$  and.



and that of the fame mulcle contracted by the power of the will, we perceive that in the mulcles there may be a flate of relaxation without their extension.

CXI. There is a flate of the contraction of mufcles that is not difpofed fpontaneoufly to alternate with relaxation, and in which too the fibres do not eafily yield to extending

powers applied: Such a flate of contraction is called a SPASM.

CXII. When muscles are excited to contraction by praeternatural caufes, and are contracted with unufual velocity and force, and especially when the contractions, alternating with relaxation, are frequently and praeter-

#### PHYSIOLOGY. 79:

praeternaturally repeated: Sush motions are called CONVULSI-ONS.

CXIII. If the contraction of mufcles are exerted with unufual force, and fuch contractions are often repeated, they, in a fhort time, become uneafy and weaker; and though contractions are not exerted with unufual force, if they are often repeated,

and for a long time, without an interval of reft, they also become uneafy and weaker.

CXIV. Within certain bounds, with refpect to force, frequency, and duration, the contraction of mufcles, by being repeated, is performed with more facility and force.

CXV.

#### So PHYSIOLOGY.

CXV. Are not the contractions of muscles produced by the action of the animal power; those which are more efpecially liable to become uneafy and weak by frequent repetition?

#### SECT. II.

CHAP. III.

OF THE FUNCTIONS OF THE BRAIN.

CXVI. From the effects of ligatures made upon the nerves, and of the destruction of their continuity, it appears, that, in their entire state, motions may be communicated from

from the brain to the other parts of the nervous fyftem, and from the latter to the former; and, from the fame experiments, it appears, that the brain (XXIX. 1.) is the organ of fenfation and volition, and of the feveral intellectual operations interveening between thefe: All which is confirmed by the effects of the organic affections of the brain upon the intellectual faculties.

CXVII. The brain is thus the fenforium or corporeal organ, more immediately connected with the mind; and, fo far as a corporeal organ is employed, all the operations of thought arifing in confequence of fenfation are operations of the brain, and are modified by its various condition; Boerb. Inft. Med. 581. Haller, Prim.



82 PHYSIOLOGY.
Prim. Lin. 570. Gaub. Path: Med.
523. See afterwards (CXXII.)

のないのないないの

CXVIII. As certain impressions act on the nervous fystem, without producing any sensation (LXXX.); and as, at the fame time, there is hardily any communication between the different parts of the nervous system, but by the intervention of the brain; it is from hence also probable, that the brain, by its organization, is fitted to propagate the motions arising in one part, to the other parts of the nervous sylftem; and, as these mechanical communications produce different effects, according to the different state of the brain itself, we, upon the whole conclude, that the brain is a corporeal organ susceptible of various con-


conditions, and thereby, of confiderable influence in most of the phaenomena of the nervous fystem.

CXIX. The action of the brain, in moving the feveral parts of the body, is excited by various caufes, or by the fame caufes in different circumftances.

1. It is especially excited by the WILL, willing the motion of certain parts as means to an end.

As the motion of certain parts is adapted to various purpofes, we are confcious of willing thefe purpofes, as they occafionally occur, and fo far alfo of the motion of the parts concerned in them; but, where the motion of the parts is connected with one fenfation, or a few only, the motions required follow thefe fenfations with-

LA CANA



#### S4 PHYSIOLOGY.


without our being conscious of specially willing them; and, unlefs we have continued the practice of adapting the motions to different purposes, we lose the power of doing fo, and the motions become unavoidably connected with those fenfations which, for a long time, had alone given occasion to them. In most of the instances of what are called VOLUNTARY MOTIONS, we are confcious of willing the end proposed, more than the motions excited; and, of the motions produced, we are confcious chiefly of those of a whole member, or of the general effect, and very little of the many particular motions that concur to produce it. We are never conscious of the particular muscles employed.

2. The

2. The action of the brain is excited by the more general and vehement volitions named EMOTIONS and PASSIONS. Upon occasion of these, the consciousness of willing the particular motions produced is always much less diffinct, and in many cafes is not at all perceived. Of the laft kind are, particularly, most of the expressions of the pasfions in the countenance and gef-

flure.

3. By the difposition of human nature to IMITATION. This imitation is sometimes involuntary, often without confciousness; and the consciousness which takes place is often of the general purpose only, without that of the particular motions produced; or, at least, it is of thefe only as a general effect. H 4. By



-4. By APPETITES or defires, directed to certain external objects, and arifing from fenfation, without any reafoning directing to an end; at leaft, without any other end in the first instance but that of the gratification of the defire.

5. By certain PROPENSITIES or defires to remove an uneafy of painful fenfation, in confequence of which motions are excited, which

are not directed to any external object, but confined to the body itself.

These motions are not forefeen; nor are we ever confcious of willing any thing but the general effect Of this kind, the chief are the motions of fneezing, coughing, fighing, hiccuping, vomiting, voiding urine and foeces, yawning, ftretching, (pandi-

(pandiculatio), and those motions of reftlefiness and inquietude which pain and uncafiness produce. Weepng and laughing are expressions of motion and passion.

In all these, as well as in the moions of No. 4. some volition is conerned, not only as they can often be prevented by another volition prefenting itself; but, besides, as the several motions which occur in

executing these propensities are more or fewer, and more or less forgible, according to the vehemence of the propensity or effort. Very often the ftimulus to these propenfities is irresisfible; and, unless the peculiar ftimulus is present, the motions cannot be produced by any volition.

H 2



6. By

### SS PHYSICLOGY.

6. By certain internal impressions arising from the exercise of the functions of the body itfelf, which produce no sensation, nor produce motions of which we are confcious, except when exercised in an unufual manner. Such are the caufes of the motions of the heart and arteries, of the organs of refpiration, of the ftomach, intestines, and perhaps of many other parts. With regard to most of these motions, it may be fupposed, that they are the mechanical effects of their feveral caufes, acting upon the inherent power of muscular fibres (LXXXVI.); but itis fufficiently certain, that they allo depend upon an action of the brain; and the effects of passions, as well as the effects of deftroying or comprefling

preffing the nerves of the organs concerned, are proofs of it. The motions mentioned in this article are commonly fupposed not to be accompanied with any volition of which we are diffinctly confcious. This perhaps is not striftly true with regard to most of them; and, so far as it is, it may be imputed to that repetition which destroys consciousness (LV.): But neither can we entirely adopt this explanation; as these motions, which are intended to follow one stimulus only, may be supposed to require no exercise of volition, as in the cafe of the heart, arteries, and alimentary canal, while the action of respiration, as adapted to various purpofes, continues to be a voluntary motion. H 3. 7. By

7. By various occafional imprefions of external bodies, and by various occafional flates of the fyftem, or of its particular parts which excite motions, not only in the parts to which the imprefions are immediately applied, but alfo in diffant parts, on which they can operate only by the intervention of the brain. Some of thefe caufes operate with, others without fenfati-

on or velition.

CXX. In all, or any of these cafes in which the action of the brain takes place, we do not perceive the manner, that is, the mechanical means, by which the several causes produce their effects; and we perceive only an institution of our Maker,

ker, establishing a connection between the feveral caules and the motions that enfue. At the fame time we, for the most part, perceive, that the connections established are fuited to the purpofe of the animal œconomy; and particularly, to the purpose of supporting the system in a certain condition for a certain time, and of averting what might hurt or destroy it. This constitution of the animal æconomy we call ATURE; and every where in the economy we perceive the vires confervatrices and medicatrices naturae, fo juftly celebrated in the schools of phyfic.

CXXI. It is in confequence of this conftitution, that not only impulfe,



pulse, and other causes which may be supposed to produce motion, do accordingly excite motions in the animal occonomy; but that also many causes, which feem to diminish motion, do however produce an increase of motion in animal bodies. Thus feveral passions which, in their first tendency, diminish motion, several propensities arising from debility and difficulty of action, the absence of usual impressions, evacutions, and other causes of relaxation, cold and narcotic powers, are all of them causes of confiderable motions arifing in the animal system.

CXXII. As the mechanism of the brain, fuited to its feveral functions, is not at all perceived (CXX.); and, at

#### PHYSIOLOGY. 93:

at the fame time, as very few of thefe functions are carried on without fenfation and volition, it must appear from this, and many other confiderations, that the mechanism of the brain (CXVII.) would not be fusicient for the purpose, without being united with a fentient principle or mind that is constantly prefent in the living fystem. But, at the fame time, it is with little probability alledged, that the administration of the corporeal functions is intirely directed by the mind acting independently of the body, and with intelligence perceiving the tendency of imprefiions, and exciting fuch motions as may favour the beneficial, or obviate the hurtful tendency, of all caufes acting upon the body. We



We are certainly confcious of no fuch administration. Many impreffions have their effects without fenfation or volition. In most cafes where volition takes place, it is very general, with little confciousness of the motions excited, and none at all of the organs employed. The force of impreffion is every where absolute; and it is according to the force of impression, and other mechanical conditions of the fystem, that the motions excited prove either falutary or pernicious. The general principle, therefore, is ill founded; it is not necessary; Vide Stabl. Praef. ad: Junker. Confp. Med.; it can be of no use, and may be hurtful to the system of physic.

CXXIII.

CXXIII. The action of the brain is often determined and regulated by cuftom and habit; that is, by laws established by frequent and uniform repetition. See above (XLIII. LIV. LV. LVI. 5. LX. 3. LXVIII. and LXX.) for the effects of cuftom on fenfation, and (CXIV.) for one effect of it on the action of moving fibres. It is now to be observed further, I. That cuftom determines the degree of tenfion(CII. CIII.) neceffary to the action of muscular fibres. 2. That custom associates motions with fenfations, which are not otherwise their causes; so that the renewal of the fensation, or of its idea, renews also the motion. 3. That cuftom affociates different motions, fo that they cannot be feparately performed, though not originally,

nally, nor necessarily, connected. 4. That cuftom determines the degree of force and velocity with which metions can be performed. 5. That -cuftom determines the order of fucceffion in affociated motions, and the velocity with which they fhall fucceed one another. 6. That cuftom citablishes the periodical return of certain feniations and motions, not originally neceffary to the æconomy. 7. That cuftom fixes an exact period for the return of certain fenfations and motions, which, by the laws of the economy, are difposed to return at intervals otherwife undetermined. Thefe laws, which may be eftablished by cuftom, are, many of them, with difficulty avoided; they are often rigidly fixed, have a confiderable influence on the



the action of the brain, and govern the revolutions of the animal fyftem. Thus, any caufes producing a deviation from the ufual degree of force and velocity (No. 4.) are apt to deftroy the measure of it altogether; and, in like manner, caufes producing a deviation from the ufual order and velocity in the fucceflion of motions (No. 5.) are apt to deftroy the power of the mind in following

that order, or in giving any meafure to the feveral motions which fhould be performed; and perhaps, in this way, the effects of debility, of feveral paffions, and of furprife, are, in fome meafure, explained.

CXXIV. The brain feems, by its conftitution, to be difposed to the alternate states of rest and activity; I as



as appears in the alternate flates of fleep and watching, which conftantly take place in every animal; but wherein this conftitution confifts, it is difficult to difcover.

CXXV. The moft common opinion is, that the brain is a fecretory organ, which fecretes a fluid neceffary to the functions of the nervous fystem; that this fluid is alter-

nately exhausted and recruited, and thereby gives occasion to the alternate flates of fleep and watching. But this supposition is attended with many difficulties. 1. It is probable, that the nervous fluid exisled in the animal embrio, before the action of the heart, or any fecretory function, could take place. 2. In animals, which, during the winter,

ter, suffer a temporary death, such' as bats, when, by heat, they are again reftored to life, the vital power of the solids is restored before the fluidity of the blood. 3. The nervous fluid fubfifts in the nerves and muscular fibres long after they are feparated from the brain, and often when cut into many fmall parts. 4. Though it be true, that the brain is: a fecretory organ, the fluid 'fecreted may be destined to another purpose; and, so far as we understand that purpose, the fluid fit for it must be unfit for the purpose of sense and motion. 5. There is no appearance, in any part of the nervous system, of provision made for an occasional accumulation of the fecreted fluid; nor is there any evidence of its actually taking place. 6. The phae-I 2 nomena



nomena of fleep and watching do not correfpond with fuch a fuppofition; as fleep often takes place, when the fecreted fluid muft be copioufly prefent, and watching can be protracted, when the fluid is exhaufted much beyond its ufual meafure. 7. Both flates are induced by many caufes, which can hardly be fuppofed to act upon a fecretion.

CXXVI. A certain compression of the brain can produce a state of the fystem refembling sleep; but that state is, in some respects, different from that of ordinary sleep: And it does not, by any means, appear, that natural and ordinary sleep depends upon any compression of the brain.

CXXVII.

CXXVII. As it is therefore probable that fleep and watching, do not depend upon a different quantity of the matter of the nervous fluid for the time present in the syflem (CXXV.), or upon any caufes interrupting its motion, while the condition of the matter remains the fame (CXXVI.); we are disposed to believe, that those states of sleep and watching depend upon the nature of the nervous fluid itself capable of becoming more or lefs moveable; that it is chiefly in the brain susceptible of these different conditions; and that, especially by its condition there, it has its more general effects on the whole system.

I 3

#### CXXVIII.



CXXVIII. This may perhaps be confirmed by confidering the remote caufes of fleep and watching : And it appears, that cold, the abfence of imprefions, attention to a fingle fenfation, or to fenfations that have no confequence in thought or action; the finished gratification of all vehement defires, fedative fensations and imprefions, evacuations, relaxation, and any violent, frequent,

or long continued exercise of the animal power, are all of them, severally or together, causes inducing sleep.

CXXIX. On the other hand, it appears, that a certain degree of heat, all fenfations of imprefion, imprefions analogous to those which produce

produce fenfation, all fenfations which lead to thought and action, and the increased impetus of the blood in the veffels of the brain, are all of them causes favouring or inducing a state of watching.

CXXX. As most of the caufes (CXXVIII.) are evidently fuch as diminish motion in the brain, and those of (CXXIX.) are such as increase it; it is from thence probable, that the nervous fluid in the brain is truly capable of different states or degrees of mobility, which we shall call its states of EXCITE-MENT and COLLAPSE; but, without intending, by these terms, to express or determine any thing with regard to the nature of the nervous



# 104 PHYSIOLOGY. nervous fluid, or wherein its diffe-

rent states confist.

CXXXI. This fubject may be further illustrated, by observing, that the excitement of the brain appears to be in very different degrees on different occasions. It seems to be greatest in certain maniacs endued with uncommon strength, resisting the force of most impressions, and

with the utmost difficulty admitting fleep.

CXXXII. A lefter degree of excitement occurs in the ordinary flate of watching in men in health, when the excitement is total with refpect to the functions of the brain, butreadily admitting of fleep. This excitement may be confidered as of two

two kinds, either as it refpects the vigour, or as it refpects the mobility of the fystem; and these different states of the brain are expressed in the body by strength or debility, alacrity or sluggissiones; and, in the mind, by courage or timidity, gaiety or fadnes.

CXXXIII. A degree of collapse takes place in the cafe of natural fleep,

when the collapse prevails fo far as to fuspend very intirely the exercise of the animal functions; and, though the exercise of the vital and natural continue, they are confiderably weakened. The partial collapse that may take place in the brain discovers itself by the delirium which appears in a state that often occurs as intermediate between fleep



fleep and watching; and even in fleep, the collapse with refpect to the animal functions takes place more or less entirely; whence the fleep is with or without dreaming, and the dreaming is more or less active.

CXXXIV. A ftill greater degree of collapfe takes place in the cafe of fvncope, in which it is fo great as to fufpend the exercise of the vital functions concerned in the circulation of the blood, notwithstanding the force of habit in these, and their being exposed to constant stimuli. Here the collapse may be very considerable; but there still remains fome degree of excitement while the brain can be acted upon by stimuli, which act only on vital powers, and

while

while its ufual excitement is still recoverable by fuch stimuli.

CXXXV. If the collapse is ftill more compleat and irrecoverable, it is the ftate of DEATH.

CXXXVI. From what is now faid of the excitement and collapse of the brain, it will appear, that we suppose LIFE, so far as it is corporeal, to confift in the excitement of the nervous fystem, and especially of the brain, which unites the different parts, and forms them into a whole. But, as certain other functions of the body are neceffary to the fupport of excitement, we thence learn that the caufes of death may be of two kinds; one that acts directly on the nervous fystem destroying its



its excitement; and another that indirectly produces the fame effect, by deftroying the organs and functions neceflary to its fupport. Of the first kind are chiefly the causes of fleep operating in a higher degree, as cold, fedative passions, poisons, and all causes of very violent excitement.

-CXXXVII. This fubject may re-

ceive ftill further illustration from confidering the flate of the other parts of the nervous fystem with respect to excitement and collapse. In the nerves strictly fo called (XXIX. 2.) we do not know that the nervous fluid suffers any change, but what is exactly correspondent to its flate in the brain and extremities; and therefore the only difference of the

the state of the nerves to be taken notice of is their being more or less free (XXX.)

CXXXVIII. In the fentient extremities of the nerves (XXIX. 3.), a difference of the flate of the nervous fluid arifes from the feveral caufes mentioned (LVI. 2. 3. 4. and 5.), which give a different degree of fenfibility; and it is probable, that thefe different flates of the fentient extremities are analogous to the different degrees of excitement in the brain.

CXXXIX. The moving extremities, or mulcular fibres (XXIX. 4.) may also be in a different condition with respect to excitement. It is probable, that their conflictution is K fuch



#### IIO PHYSIOLOGY.

fuch as to admit of a higher degree of excitement than any other portion of the nervous fystem; and that upon this their contractility depends But, whatever is in this, we perceive very clearly, that the condition of muscular fibres may be varied by caufes affecting their tonic power (CI.), or their vigour and mobility (LXXXIX.), and by the effect of cuftom (CXIV.); and it is probabl, that the states produced by thefe causes are analogous to the different degrees of excitement in the brain (CXXX.), and in the fentient extremities (CXXXVIII.); and thus the feveral parts of the nervous fyftem (XXIX.), as they confift of the fame kind of matter (XCIV.), are also subject to similar conditions.

CXL

CXL. The beginning of motion in the nervous system is most commonly accompanied with fenfation, and the force of this in producing its several effects is more or less, I. According to the force (XLII.), quality (LXXVII.), and novelty (LIV.) of imprefion. 2. According to the fenfibility of the fentient extremity and brain (LVI.). 3. According to the ftate of attention (XLVII.). These several causes often concur, frequently balance one another, and are always to be confidered together.

CXLI. The effect of fentation is commonly that of exciting the action of the brain; and this is generally according to the degree of voli- $K_2$  tion



112 PHYSIOLOGY. tion produced under the different circumftances mentioned in (CXIX.)

CXLII. The action of the brain excited by volition or other caufes, is determined to particular parts of the body, most constantly by the connections established in the system (CXX.), but also occasionally, by acquired habits, or the greater mo-

# bility of certain parts.

CXLIII. With regard to the connections eftablished in the fystem (CXX. CXXI.), it is to be observed, as of great confequence in pathology, that certain parts of the body which have a common function and conftitution, have thereby a peculiar relation to the brain, so as to be more liable to be affected by the different

different flates of it, and in their turn by the difference of their condition to affect the brain. Such are, efpecially the organs of voluntary motion; the alimentary canal, and efpecially the flomach; the circulatory fyftem, and particularly the extremities of the veffels on the furface of the body; the uterine and genital fyftem in females; and fome others to be mentioned in the pathology.

CXLIV. The communications of motion between the feveral parts of the nervous fyftem which have been mentioned as inftances of a particular fympathy between these parts, are very feldom to be explained by any contiguity or contact, either in K 3 the




the origin or courfe of the nerves of the communicating parts; and more commonly they may be explained, by fuppofing the action of the imprefion to be general with refpect to the brain; and that the affection of particular parts depends upon the caufes of determination (CXLII. CXLIII.)


When the action of feveral parts, together or fucceffively, are neceffary

to produce one effect, these concur, though the stimulus exciting the action of the brain is applied to one fingle part only; and commonly no other cause of communication can be assigned, but the several motions being necessary to the execution of the volition, propensity, &c. arising from the stimulus.

CXLV.

CXLV. Thefe are the chief facts and laws relative to the nervous fyftem. The whole might perhaps be illuftrated, and more exactly afcertained by a more particular inquiry into the nature of the nervous fluid; but we are not fo confident in our opinion on this fubject, or of the application it will admit of, as to deliver it here.





# ΡΗΎSΙΌLΌĠΎ.

#### SECT. ΠI.

OF THE CIRCULATION OF THE BLOOD.

CXLVI. The circulation of the blood, by (CXXIX.), feems neceffary to the excitement of the brain; and for this, as well as other reafons, it is, next after the function of the brain itself, the most important of the animal œconomy.

On this fubject, we shall confider, 1. The course of the blood. 2. The powers by which the blood 15 moved.

. . .

moved. 3. The laws and general circumftances of the circulation.
4. Refpiration as neceffary to it. On all these fubejcts, we fuppose the anatomy of the parts to be known.

 $S \in C T$ . III.

#### CHAP. I.

OF THE COURSE OF THE BLOOD.

CXLVII. Wounds and haemorrhagies fhew, that, in living bodies, the blood is in conftant motion, and flowing into any one part from many others.

CXLVIII.

### IS PHYSIOLOGY,

CXLVIII. In man, and other analogous animals which have once breathed for fome time, the courfe of the blood is very conftantly in the following manner.

From the left ventricle of the heart, the blood palles into the trunk of the aorta, and fucceffively into the fellowing veffels and cavities, viz. The branches of the aorta, the branches of the *vena cava*,

the trunk of the fame, the right auricle of the heart, the right ventricle of the fame, the pulmonary artery, the pulmonary veins, the left auricle of the heart; from which laft it paffes into the left ventricle of the heart to return again into the fame courfe as before.

From all this, it appears that, in the arteries, the usual course of the

the blood is from the heart, towards the extreme branches of thefe; and that in the veins, the courfe is in the contrary direction, from the extreme branches towards the heart.

CXLIX. The course of the blood through the cavities of the heart, as above described, is not in a continued ftream, but alternately interrupted and free during the contraction and dilatation of these cavities, which alternately happen. Thus, while the left ventricle of the heart is in a state of contraction, the blood passes out of it into the aorta; but at the same time no blood passes into it from the left auricle, which is then dilated and filled by the blood flowing into it from the pulmonary vein. It is only when the ventricle is emptied .

•

emptied by contraction, and confequently relaxed, that the blood passes into it from the auricle, urged by the contractions of the auricle and adjoining sinus venosus, which fucceed immediately to that of the ventricle. During this contraction of the auricle and filling of the ventricle, no blood passes from the ventricle into the aorta, nor till a contraction of the ventricle succeeds in consequence of its being fil-The same circumstances take led. place with regard to the right ventricle and auricle of the heart, and precifely at the fame times; for it appears, that the two ventricles d the heart are contracted and relaxed at the fame time; and, in like manner, the two auricles.

CL. That the course of the blood, as defcribed (CXLVIII. and CXLIX.), is its usual and constant course, appears from the infpection of the heart in living animals; from the fituation of the valves of the heart; from the fituation of the valves at the orifices of the aorta and pulmonary artery; from the fituation of the valves of the veins; from the effects of ligatures made upon the arteries and veins; from the effects of haemorrhagies of the veins; from obfervations with the microfcope; and, lafily, from experiments of infusion and transfusion in living, and of injection in dead animals.

L

CLI.

CLI. This course of the blood is, however, in some parts, and upon some occasions, a little changed. I. In the penis, and fome other parts, the blood does not pass from the extreme arteries immediately into continuous veins, but is poured out into an intermediate cellular texture, from which it is afterwards received by the extreme veins. 2. In the small branches of the aorta, the blood does not move constantly in a direction from the heart towards the extremities; but, in certain portions or them, is sometimes retrograde to that courfe. In this, it is favoured by the anaflomofes frequent between the finall veffels, which, at the fame time, prevent this deviation from being confiderable

PHYSIOLOGY. 123. ble or durable, while the action of the heart continues.

CLII. In the course of the venous blood, there is also some variety. 1. In the fmall veins, the blood is liable to have a motion retrograde to its usual direction (CXLVIII.), as in the arteries (CLI. 2.) 2. The blood returning to the heart from moft parts of the body, passes fuccessively from sinaller into larger veins, by a feries of pretty regularly increasing vessels, till they form the vena cava entering the heart. But this is varied in the abdomen, where the veins carrying the blood returning from every vifcus included in that cavity, except the kidneys and genitals, unite in forming the vena: L 2 portarum,

portarum, whereby they undergo a peculiar distribution. 3. The veins returning the blood from the extreme arteries in the brain do not carry it to the heart by a series of regularly increasing vessels, but by the interposition of finuses, into which the fmall veins iffuing from the cortical part of the brain immediately pour their blood. 4. As the course of the blood through the veffels of the lungs is not at all times equally free, and particularly, is confiderably interrupted at the end of expiration; so at the same time it is alfo interrupted at entering the right ventricle of the heart; and this often occasions some regurgitation, or retrograde motion, in both the afcending and defcending cava.

CLIII.

CLIII. The whole of the fluids carried in the aorta to its extreme branches, are not returned again by the continuous veins to the heart, as, by fecretory veffels, a part of them is conftantly carried out of the course above described. Some of these secreted fluids are thrown intirely out of the body, and others are poured into certain cavities, for various purposes of the æconomy. Of these last, there is a peculiar fluid which, from the extremities of the arteries, is poured out in a liquid form, or exhaled in that of vapour, into, perhaps, every cavity and vacuity of the body. This, after having ferved the purpose of the effusion, seems intended to be regularly returned again into the course of the circulation; and accordingly, in all the fe- $L_3$ veral

veral cavities into which it had been effused, there are absorbents which again take it in. These do not carry the fluid immediately into the veins, but uniting form the veilels called LYMPHATICS, which, in their course, pass through the conglobate glands, and at length terminate either in the receptacle of the chyle, in the thoracic duct, or in the left subclavian vein; and, in this way, return the absorbed fluid into the course of the ordinary circulation.

CLIV. There are abforbent veffels, not only in all the feveral cavities, but alfo on the external furface of the body, by which many extraneous

# PHYSIOLOGY. 127 extraneous matters may be introduced into it.

CLV. Moft of the fluids fecreted from the circulating mafs, and poured into cavities, may be abforbed from thefe, and returnedagain by the lymphatics, as in (CLIII.), to the courfe of the circulation. But the fame fecreted fluids feem often to be returned alfo into the courfe of the circulation by a regurgitation, or retrograde motion, in the excretory and fecretory veffels.

#### SECT.

#### SECT. III.

#### CHAP. II.

## OF THE POWERS MOVING THE BLOOD.

CLVI. The chief power by which the blood is moved, and the circulation carried on, is the action of the heart, or its repeated contractions occurring, as mentioned (CXLIX.) For this purpofe, the heart is a mufcular part; the action of which may depend upon an inherent power ftimulated by the dilatation of its cavities;

vitics; but this inherent power requires the conflant fupport of the nervous and animal powers; and is often actuated by thefe.

CLVII. The contraction and relaxation of the heart; or, as thefe are called, its fyftole and diaftole, are neceffarily alternate by the general law (CVIII.) affecting all mufcles; and by the ftimulus from the influx of venous blood being alternately applied and removed.

CLVIII. If we may be allowed to eftimate the vigour of muscles by the number of their fibres, we must suppose the force of the heart to be very confiderable; but it is very difficult to obtain any exact eftimate

of

r30 PHYSIOLOGY. of its absolute force. It is perhaps the relative force only that we are concerned to know.

CLIX. Do the arteries; by their contraction, contribute to promote the motion of the blood begun by the heart? They cannot do it by the elasticity of the fimple solid in their structure, and can do it only by their being endued with a muscular power, whereby they may, in their contraction, communicate to the moving blood more force than was lost of the force of the heart in dilating them. That they are indued with a muscular contractility (LXXXII.), is probable from the appearance of the muscular fibres in their structure; from their irritability appearing in the experiments of Verschuir; from their becoming flaccid

cid on tying the nerves belonging to them; from the motion of the blood being fupported, when the force of the heart is confiderably weakened; from the motion of the blood becoming languid, when the action of the arteries is deftroyed; from the velocity of the blood in the extreme arteries being greater than was to be expected from the velocity of the blood iffuing out of the heart, exposed to fo many causes of retardation as constantly occur; and, laftly, from the velocity and impetus of the blood in different parts of the body, and at different times, being unequal, while the action of the heart continues the fame.

It is probable, that the muscular fibres of the arteries become more irritable

32 PHYSIOLOGY. ritable as the arteries are more diffant from the heart.

CLX. The tone and action of the arteries, as a mufcular part, may be increafed by ftimuli immediately applied to them, or by the increafed force of the nervous and animal powers with regard to them; and they may be diminished by fedative powers applied, or by weakening the nervous and animal powers.

CLXI. There does not appear to be any ofcillatory motion in the extreme arteries independent of the action of the heart.

CLXII. There does not appear to be any operation of capillary attraction

traction in the extreme arteries; nor does there feem to be any occasion for fuch a power in any part of the arterial fystem.

CLXIII. The power of derivation (Vis derivationis Ill. Halleri) in the fanguiferous fystem, seems to be no other than that which arises from the fulness of contractile veffels.

......

CLXIV. The motion of the blood in the arteries of any particular part, is promoted by the action of adjoining muscles.

CLXV. The blood in the vena cava, and its branches, is moved by the action of the heart and of the arteries. These powers are affisted M by

by the action of mufcles, which, in their contraction, prefs the veins lying between their feveral fibres; and alfo, by the fwelling of their whole mais, prefs the adjoining veins. Thefe veins are commonly provided with valves, which determine the effect of all prefiure upon them, to be the motion of the blood towards the heart.

The great trunks, both of the

vena cava and pulmonary vein, are provided with mulcular fibres, and manifely endued with mulcular contractility.

CLXVI. In the abforbent veffels, the fluids are probably taken in by a capiliary attraction.

CLXVII.

CLXVII. In the lymphatic veffels, provided with numerous valves, which neceffarily determine the motion of the contained fluid to be towards the heart, the fluid is moved by the preffure of the neighbouring mufcles and arteries. But further, as the lymphatics are remarkably irritable, it is probable that the fluid in them is moved by a periftaltic motion begun by the action of their

absorbent extremities.

CLXVIII. The motion of the blood through the veffels of the lungs, depends upon respiration, to be confidered hereafter.

 $M \cdot 2$ 

SECT.

# SECT. III.

#### CHAP. III.

# OF THE LAWS OF THE CIRCULA-TION.

CLXIX. The velocity of the blood paffing out of the left ventricle of the heart into the aorta, may be eftimated from knowing the quantity of blood paffing out at each fyftole, the area of the orifice of the aorta, and the time occupied by the fyftole; but none of thefe circumftances are exactly afcertained. CLXX.

CLXX. As the blood moves onwards through the arteries, the velocity (CLXIX.) fuffers a confiderable retardation, from several causes. I. From the capacity of the arterics being enlarged as they are more difant from the heart. 2. From the frequent flexures of the arteries. 3. From the angles which the branches. make with the trunks from which they arife. 4. From anastomoses. 5. From the viscidity of the blood. 6. From the friction of adhesion. 7. From the weight and rigidity of the parts furrounding the arteries.

CLXXI. The velocity (CLXIX.), and the caufes of retardation (CLXX.) being given, the velocity of the blood in the arterics will be as the M.3 frequency



# 138 PHYSIOLOGY. frequency of the fystole of the heart.

CLXXII. The frequency of the fyftole of the heart will be more or lefs, 1. As the blood in the veins is more or lefs quickly returned to either ventricle of the heart. 2. As the ventricles of the heart are more or lefs intirely evacuated at each fyffole: 3. As the mulcular fibres of the heart are more or lefs moveable. 4. As the action of the nervous and animal powers are more or lefs increafed with refpect to the heart.

CLXXIII. As the arteries of a healthy body are always full, the blood thrown out of the ventricles into the arteries during the fystole of the heart, can only find a place there by

by pufhing on the blood with the velocity (CLXIX.), or by dilating the arteries; but as the refiftances (CLXX.) prevent the blood fromflowing with the velocity (CLXIX.), the blood thrown out of the heart muft, in fome measure, dilate the arteries, and thereby form what is called the PULSE.

CLXXIV. It appears, that, in the

arteries, to a certain length, the blood moves fafter during the fyftole than during the diaftole of the heart; but, as the refiftances and caufes of retardation become greater in every portion of the arteries, as it is more diftant from the heart, fo the acceleration of the blood during the fyftole of the heart muft be greater in any portion of the arteries nearer the heart



heart than in the next adjoining; that is more diftant; and, fo far as this takes place, a dilatation of the arteries will happen, even from a fmall quantity of blood thrown out of the ventricles.

CLXXV. As the refiftances to' the blood's motion in the bloodveffels increafe with the diftance from the heart, there may be a part of the fanguiferous fyftem in which' the motion of the blood will not be accelerated during the fyftole of the heart, and in which, therefore, no pulfe can be difcerned. This happens in the extreme branches of the aorta; and no pulfe is ever obferved in the extreme branches of the vena cava.

CLXXVI.

# PHYSIOLOGY. 14T

CLXXVI. The velocity and impetus of the blood in the whole fyftem of blood-veffels will always be as the action of the heart and arteries taken together.

CLXXVII. The velocity and impetus of the blood in any particular part of the fyftem will be, 1. As the part is more or lefs diftant from the heart. 2. As the circumftances (CLXX.) take place more or lefs in the part. 3. As the gravity of the blood concurs with, or oppofes its motion in the part. 4. As caufes increasing or diminishing the action of the arteries of the part are applied or removed.

#### CLXXVIII.

.

CLXXVIII. The quantity of blood diffributed to any particular part of the fanguiferous fyftem, will be greater or lefs according to the velocity and impetus of the blood in the part, by (CLXXVII.); and according to the refiftances in other parts being increafed or diminifhed by confiriction, compression, ligature, position, relaxation, or aperture.

CLXXIX. The flexibility and contractility of the blood-veffels render the effects of all increase or diminution of refistance in any particular part most confiderable in the nearest, and very little fo in the more remote veffels of the fystem. By this we are to judge of the celebrated

PHYSIOLOGY. 143 lebrated doctrines of derivation and revullion.

CLXXX. The quantity of blood diffributed to the different parts of the fyftem, is in different proportion at different periods of life.

I. The capacity and force of the heart, in proportion to the fyftem of veffels, is greater at the beginning of life than at any after period. Till the body arrives at its full growth, the capacity of the veffels increafes in greater proportion than that of the heart; but, from that period, the capacity of the veffels is conftantly diminifning, while that of the heart fuffers little change.

2. A greater quantity of blood is contained in the arteries, in proportion to that which is contained in the veins,

veins, at the beginning of life, than at any after period. From the time that the body has arrived at its full growth, the quantity of blood contained in the veins, in proportion to that which is contained in the arteries, is conftantly increasing.

3. The veffels of the head receive a greater quantity of blood in proportion to the reft of the fyftem, at the beginning of life, than at any after period.

4. Any general increase of the action of the heart and arteries, determines the blood more copiously to the extreme arteries on the furface of the body, than to those of the internal parts.

5. The equilibrium of the fanguiferous fystem, with regard to the distribution of the blood, may be

be changed by various caufes. (CLXXVII. CLXXVIII.); and thefe. caufes continuing to operate for fometime, induce a habit which renders the changed diffribution neceffary to the health of the fyftem.

6. The lymphatic fystem is fuller in young perfons than in old.

# S E C T. III.

# CHAP. IV. Of Respiration.

CLXXXI. Refpiration confifts of the motion of infpiration, or the admiffion of air into the lungs, and of expiration, or the expulsion of N air

# 146 PHYSIOLOGY. air from the fame, alternately happening.

CLXXXII. Refpiration takes place in man, and other analogous animals, foon after the infant is taken from the uterus of the mother, and is exposed to the air. After it has taken place for a little time, it is ever after neceffary to the continuance of life, as it is absolutely neceffary

to the continuance of the circulation of the blood.

CLXXXIII. The lungs are a hollow fpungy mafs, capable of confining air, and readily dilatable by it. By the wind-pipe, they are open to the atmosphere; and they are fo fituated in the thorax, that the air must enter into them, if the cavities of

of the thorax, in which they are placed, are enlarged. For, as there is no air in these cavities, and the external air cannot enter into them, the enlargement of the thorax must form a vacuum around the lungs, which the external heavy and elaflic air will supply by entering into and dilating the lungs, while these do not allow the air to pass through them into the cavities of the tho-

rax.

CLXXXIV. Infpiration therefore depends upon the enlargement of the capacity of the thorax; and this is performed chiefly by the contraction of the diaphragm. This, in its relaxed ftate, is fufpended by the mediaftinum, and its middle tendinous part is raifed high in the N 2 thorax

thorax; wherefore, as this middle part, by the contraction of the mufcular, is moved downwards, the thorax is thereby confiderably enlarged.

CLXXXV. The capacity of the thorax is also enlarged by the motion of the ribs upwards, whereby the curvatures of opposite ribs are fet at a greater diffance from each o-

ther; and, by the fame motion, the flernum is moved outwards, and fet at a greater diffance from the vertebrae of the back. The motion of the ribs upwards is caufed by the contraction of both layers of intercoftal mufcles. That the mufcles called internal intercoftals concur with the external in raifing the ribs, appears from the fituation of thofe mufcles,

mulcles, from the greater mobility of the inferior ribs, from the infpection of thole mulcles in living animals, and from experiments imitating their action. In more violent and laborious infpirations, the raifing of the ribs is affifted by many mulcles attached to the ribs, and arifing from the clavicle, humerus, fcapula, and vertebrae of the neck or back.

CLXXXVI. By the enlargement of the thorax, a dilatation of the lungs is produced, in proportion to the bulk of air entering into them; but the dilatation may often be greater by the air that enters into the lungs being heated and rarified; and the greateft diftenfion of the lungs is N 3 obtained

obtained by a confiriction of the glottis confining the air that has already entered into the lungs.

CLXXXVII. As infpiration, or the admission of air into the lungs, depends upon the enlargement of the thorax, the diminution of it must expel the air, or produce expiration. The capacity of the thorax is diminished, while the muscles dilating it are fpontaneoufly relaxed, by the elafficity of the ligaments connecting the ribs with the vertebrae, and by the elafticity of the cartilages and ligaments connecting the ribs with the fternum; both which powers, commonly affilted by the weight of the ribs themfelves, bring the ribs and sternum into the position they were

were in before inspiration. At the fame time, the elasticity of the mediaftinum draws the diaphragm upwards; and the contraction of the abdominal muscles both prefies the diaphragm upwards, and pulls the ribs downwards; and, in the last, they are affifted by the sterno-costal and infra-costal muscles. While these powers concur in diminishing the capacity of the thorax, the expulfion of the air from the lungs is affifted by the elafticity of the lungs themselves, and by the contraction of the mulcular fibres of the bronchiae.

CLXXXVIII. Thefe are the ordinary powers of expiration, which, depending upon the reaction of elaftic

#### FJZ PHYSIOLOGY.

laftic parts, is performed flowly, and with little force; but, when it is neceffary to perform it with more velocity and force, fome other and very powerful mufcles, as the quadratus lumborum, facrolumbalis, and longifimus dorfi, concur in pulling down the ribs; and, at the fame time, the abdominal mufcles, actuated by the animal power, are contracted with greater velocity and

force than in fpontaneous expiration.

CLXXXIX. The fituation of the blood-veffels of the lungs is fuch, that, in the contracted ftate of this vifcus, thefe veffels muft be much folded and ftraitened; and it appears, that, in the fœtus where they are conftantly

conftantly in a contracted ftate, their capacity is not fufficient to tranfmit, in the time required, the whole of the blood returning to the heart by the vena cava; but, after refpiration has been repeated for fome time by the dilatation of the lungs to a certain degree in infpiration, their blood-veffels are unfolded, lengthened, and enlarged, fo as to be capable of tranfmitting the whole blood of the cava.

CXC. In the infant who has breathed for fome time, the whole blood of the *vena cava* paffes into the right ventricle of the heart, and from thence enters into the veffels of the lungs; but, in the contracted ftate of the lungs, which occurs at the end of expiration, the blood can-

not

not be properly transmitted; and, for that purpose, an inspiration becomes absolutely necessary.

CXCI. It is, however, under a certain degree of infpiration only, that the blood is freely transmitted through the veffels of the lungs; for, if the infpiration is full, and continued, fo that the lungs are thereby much diffended, we find that this flate alf interrupts the free paffage of the blood, and renders expiration necessary. Expiration becomes also neceffary; because, perhaps, the air long retained in the lungs lofes a part of its elasticity, and becomes thereby unfit to keep the lungs diftended; but, more certainly, and more efpecially, because, in an animal that has breathed

breathed for fome time, there is a noxious vapour conftantly arifing from the lungs, which, if not diflolved by the air, and carried out of the lungs, proves pernicious to life.

CXCII. From what has ben faid, it appears, that the alternate motions of infpiration and expiration are neceffary to the circulation of the blood, and otherwife, alfo to the blood, and otherwife, alfo to the health of the body; and it appears alfo, that the more frequent the alternate motions of refpiration are, the more quickly is the blood tranfmitted from the right to the left ventricle of the heart.

CXCIII. We can now perceive alfo the caufes exciting these alternate motions;

mitions; and we find no occasion for fuppoling them to arife from any caufes alternately interrupting the motions of the nervous fluid, or of the blood into the muscles concerned in these functions.

Infpiration, or the action of the muscles producing it, is excited, in all cafes of general effort, to remove pain and uneafinefs; and it is, perhaps, a propenfity of this kind that gives the first beginning to refpiration in the new born infant, exposed to several new and uneasy imprefiions. For the continuance of respiration, infpiration is efpecially excited by the fenfe of uncafinels that attends any difficulty in the paffage of the blood through the veffels of the lungs; but this uneafinefs arifes, in fome

## PHYSIOLOGY. 157 fome measure, at the end of every expiration, and is much increased by any continuance of this state.

CXCIV. Expiration, in fome measure, necessarily succeeds inspiration, by the fpontaneous relaxation of the infpiratory muscles (CVIII.), while the elafticity of the membranes, ligaments, and cartilages ftretched in infpiration, brings back the ribs and diaphragm into their former fituations; and the fame effects are also produced by the action of the abdominal muscles, and of the muscular fibres of the bronchia; both of which are firetched, and thereby excited in infpiration.

In the cafe of ordinary infpiration, these causes are sufficient to produce a spontaneous expiration. But, O . as

as it appears, that a violent and long continued infpiration interrupts the paffage of the blood through the lungs, this creates an uneafinefs, and a propenfity, which must produce a relaxation of the infpiratory, and excite a contraction of the expiratory muscles.

It is further to be fuppofed, that, in animals which have breathed for fome time, cuftom has affociated the feveral motions concerned both in infpiration and expiration; fo that an irritation applied to any part of them neceffarily excites the whole; and it may alfo be fuppofed, that habit determines these motions regularly to fucceed one another.

CXCV. In this manner (CXCIII. CXCIV.) refpiration is continued for

for the general purposes of the animal œconomy; but the feveral motions of which it confifts are also occafionally excited, and varioufly modified by the will, intending particular effects to be produced by these motions. They are also excited, and varioufly modified by certain emotions and paffions, and give particular expressions of these. They are often excited also by imitations; and they are particularly excited by propensities to remove pain and uneasiness, which operate more frequently on respiration than upon any other function.

CXCVI. The confideration of the effects of refpiration on the animal fluids, is delayed till the nature of these fluids shall have been more generally confidered.

02 . PHY-

#### SECT. IV.

#### OF THE NATURAL FUNCTIONS.

CXCVII. The animal body, from a fmall beginning, grows to a confiderable fize, and at the fame time, from the period of the birth, during the whole of after life, the body fuffers, by various means, a daily and confiderable wafte.

CXCVIII. The increase of bulk, therefore, must be acquired, and the daily waste supplied, by matters taken into the body, the most part of which,

which, from the prefumed purpofe of them, we name ALIMENTS.

CXCIX. A great part of these aliments, as taken into the body, are of a different nature from the matter of the body itself, or at least, are in fuch a state as not to be fit for being immediately applied to the purposes of it; they must, therefore, be changed, and sitted to the purposes of the œconomy by powers within the body itself.

CC. The conversion, or affimilation of the aliments to the nature of the folids and fluids of the animal body; the farther changes of these fluids, for various purposes, by fecretion ; and the application of some part of  $O_3$  them

them in nutrition, or in increasing the growth of the body; make what are called the NATURAL FUNC-TIONS.

#### SECT. IV.

CHAP.I.

#### V ++ ++ ++

#### OF DIGESTION.

CCI. The term *digeftion* is commonly employed to fignify the function of the ftomach alone in changing the aliments; but, in this chapter,



ter, we are to confider all the changes of these as they occur successively in the different stages through which the matters pass.

CCII. Animals are determined to take in aliment, by the appetites of hunger and thirst.

CCHI. Hunger is an appetite de-

pending upon a fenfation referred to the flomach, and arifing from a particular flate of it.

This flate fcems to be in fome refpect the degree of emptinefs, but more efpecially, the flate of contraction in the mufcular fibres which emptinefs gives occafion to. This flate of contraction may also be excited by certain flimulants applied ; but,

•

but, more commonly, it depends upon, and is correspondent to the flate of inanition, and therefore of contraction, in the vefiels of the fkin emitting the matter of perspiration.

CCIV. Thirft is an appetite for liquids, which depends upon a fenfation chiefly referred to the internal fauces, and arifing from the drynels or heat of thefe parts ; from acrimony applied to them, or exifting in the fluids poured out there; from the putrefcency or vifcidity of the contents of the flomach; and from all increafed evacuations.

CCV. These appetites determine men to take in a great variety of solid and liquid matters, directed by inftinctive

## PHYSIOLOGY. r65 inftinctive likings and difgufts; in fome inftances corrected by experience.

CCVI. Of the matters chofen, it appears that fome of them are fuited to fupply the matter of the fluids or folids of the body, and, therefore, properly named Aliment ; while others of them are fuited only to improve the relifh of aliment, or to obviate fome deviations ready to happen in the bufinefs of digeftion; and thefe we name CONDIMENTS.

CCVII. The proper alimentary matters are animal or vegetable only.

CCVII.

CCVIII. The animal aliments feem to be fo nearly of the fame nature with the matter of the body itfelf, that, to be rendered fit for the purposes of the œconomy, they seem to require no other change, but that of being rendered fluid.

CCIX. But the vegetable aliment is very different from the matter of

the animal fluids or folids, and muft therefore be changed into the nature of the fe by the powers (CXCIX.); and, as many animals are nourifhed by vegetable aliment alone, and as, perhaps, all animal matters may be ultimately traced to a vegetable origin, it will appear, that, to account for the production of animal matters, it is effectively, and in the first place,

place, neceffary to fhow how vegetable matter may be converted into animal.

CCX. If we confider the many differentodors, taftes, and colours, which are to be observed in different vegetables, we should be ready to think that vegetable matter is of very great variety; but we know that the matter diftinguished by its sensible qualities makes but a small part of the whole of any vegetable, and that, befides the matter peculiar to each, there is in most, perhaps in all, vegctables, a large proportion of common matter, which we prefume to be the matter adapted, and that very universally, to the aliment of animals.

CCXI. .

CCXI. It is this common matter of vegetables, therefore, that we are to confider here; and we think it may be confidered as of three kinds only, that is, oily, faccharine, and what feems to be a combination of thefe two.

CCXII. The oily matter-of vegetables, which makes part of the aliment of animals, is without any fenfible odor or tafte, and is not only very nearly the fame in the many different vegetables from which we take it, but is alfo in all of thefe fo nearly a kin to the oil which appears in animals, that it is not neceifary to fuppofe any confiderable change to be made upon the vegetable

table oil on its being taken into the bodies of animals.

CCXIII. It is the faccharine matter, and efpecially this when blended with oily matter in different proportion, that makes the greateft part of the common matter of vegetables, and is the chief part of the vegetable aliment of animals. It is this, therefore, that we have efpecially to confider here; and, as it lies in vegetables, it is different from the most part of animal matters in the following respects.

It is readily fulceptible of a vinous and acetous fermentation, and fpontaneoully enters into the one or the other of these; and, without undergoing more or less of these, it, P perhaps,

perhaps, never enters into a putrefactive fermentation.

The fame matter treated by diffillation, without addition, gives out always, in the first part of the diffillation, an acid, and only afterwards a volatile alkali in finall proportion.

The fame vegetable matter, treated by calcination, leaves afhes, which contain a fixed alkali, and an earth that is or may be converted into a quick-lime.

CCXIV. In all these respects, the common matter of animals is considerably different.

This enters fpontaneoufly into a putrefactive fermentation, and that without paffing through the vinous

or

•

## PHYSIOLOGY. 17r

or acetous: At least, these are not to be diffinctly perceived.

The fame animal matter, treated by diftillation, gives out always, in the firft part of the diftillation, a volatile alkali in large proportion, and only afterwards, by a great force of fire, it gives out an acid.

Animal matters, treated by calcination, leave afhes, in which no alkali is to be found, and the earth is

not calcarious, nor convertible into a quick-lime, by any means yet known.

CCXV. These differences are fufficiently marked; but it is proper to observe here, that the vegetable matter we treat of, by undergoing a putrefactive fermentation, is chan-P = 2 ged

ged fo, as to acquire very exactly most of the characters of animal matter we have just now mentioned.

CCXVI. The aliment being thus confidered, we proceed to confider the changes it undergoes after being taken into the animal body; but, firft, fof the courfe it paffes through, and of the motions it is fubjected

to in its progress.

CCXVII. The aliment is taken into the mouth, and there the more folid parts of it are commonly fubjected to a triture, or what is called manducation. At the fame time a quantity of faliva, and of the other fluids of the mouth, with fome portion of our drink, is intimately mixcd

ed with it, whereby the whole is reduced to a foft pulpy mafs. In this flate, by the action of deglutition, it paffes through the fauces into the oefophagus, by which it is conveyed into the fiomach.

CCXVIII. Here the aliments is detained for fome time, fubjected to a conftant agitation and fome preffure, both by the contractions of the different parts of the ftomach itfelf, and by the alternate preffure of the diaphragm and abdominal mufcles. After fome time, however, first the more fluid parts, and, at length, the most minute parts of the folid matter are pushed through the pylorus into the duodenum.

CCXIX. The matters received from the flomach into the duodenum pafs P 3 on

on from thence fucceflively through the feveral parts of the inteffinal canal; and, in the whole of the courfe, the matters are ftill fubjected to the alternate preflure of the diaphragm and abdominal mufcles, and to the contractions of the inteftincs themfelves.

CCXX. Through the whole course of the inteffines, but especial-

ly in thefe named *the fmall*, the more fluid part of the contents, and particularly the peculiar fluid we name chyle, is taken into the veifels named *lacteals*. Thefe, from imperceptible beginnings on the internal furface of the inteftines, unite into larger veffels laid in the mefentery, and convey the chyle, and what accompanies it, first into the conglobate glands

glands of the melentery, and from thence to the *receptaculum chyli*, as it is called. From this the chyle pafles by the thoracic duct into the left fubclavian vein. In one or other part of this courfe of the chyle, the veffels carryng it are joined by lymphatics, returning the lymph from almost every part of the body.

CCXXI. The matters contained

in the inteffinal canal, not taken into the lacteals, are moved onwards in the courfe of the inteffines, becoming by degrees of a thicker conliftence, efpecially in the colon, where their motion is confiderably retarded; but, at length, they are moved onwards to the extremity of the rectum, where their weight; bulk, and acrimony excite motions which

## 176 PHYSIOLOGY. which throw them entirely out of the body.

CCXXII. This is the course of the alimentary matters, so far as they can be considered as any ways in a separate state. Of the motions of the feveral organs concerned in this course, we pass over those of manducation, deglutition, or others depending on the action of muscles, the

functions of which are readily underftood from a knowledge of their fituation; and we are here to confider only the motions of the alimentary canal itfelf.

CCXXIII. The motions in the oclophagus depend upon the action of its mulcular fibres, which are chiefly those forming a chain and circularly



larly furrounding it. This tube, by the morfel of food puflied into it by the action of deglutition, is neceffarily dilated, and its circular fibres are thereby excited to a contraction. But as these fibres are fucceffively dilated, so are they also contracted and push on their contents through the feveral portions of the tube, alternately and fucceffively dilated and contracted, giving the appearance of a vermicular motion and what is commonly called periftaltic. This motion may be propagated either upwards or downwards, and the direction of it is in the one or the other way, as the motion happens to begin at the upper or lower extremity.

CCXXIV. The motion of the flomach is not fo fimple. Its mulcular fibres

fibres are in like manner irritable by dilatation, and its circular fibres must therefore be in some measure subjected to a fucceffive dilatation and contraction. But, though the direction of fuch motions is from the left to the right, this does not immediately push the contents of the ftomach into the intestines. It seems to be the purpole of the æconomy, to detain the aliment for some time in the itomach; and therefore, any confiderable dilatation of the circular fibres, especially that which occurs in a full ftomach, feems to have the effect of exciting the longitudinal fibres to a contraction, which draws the two orifices of the ftomach nearer to one another. By this the pylorus is raifed up and rendered less eafily passable, and probably,

bably, at the fame time, the peculiar band of circular fibres which furround the pylorus, are more firmly contracted, and render it less pervious. Vide (CXLIV.) From hence it is, that the direction of the peristaltic motion of the stomach is fometimes from the left to the right, and fometimes also the contrary way. It is, however, most constantly in the first manner; because it is commonly begun from the œfophagus, and becaufe, when it is inverted, the refiftances on the left from the blind fack of the ftomach, from the higher fituation of the cardia, and from the conftriction of this by the diaphragm in infpiration, are commonly more confiderable than the relistance at the pylorus. The contents of the flomach, therefore, are at

at length pushed through the pylorus; in the first place, the more fluid contents, as these occupy the antrum pylori; while the more folid, having their air loofened by fermentation, are rendered specifically lighter, and float nearer the upper orifice. But at length, as the ftomach is in any measure emptied, the pylorus is less raised, is more relaxed, and allows matters to pass more eafily; and, at the fame time, the empty stomach contracted more, is eipecially towards the right extremity contracted to fuch a degree, as to embrace the smallest solid matters, now fallen down into it, and to puth them through the pylorus. This is an idea of the ordinary motions of the flomach; but they are, upon some occasions, subject to other modi-

modifications, as in eructation, rumination, and vomiting, which, however, as morbid, we referve to be confidered in the pathology.

CCXXV. The motions of theintestines will be readily understood, from what has been faid of those of the oefophagus. Any portion of the intestinal canal being dilated, will in consequence be contracted, and will urge on its contents in the fame direction in which the motion was begun. But as the force here is gentle, and as, in the long course of the canal, there occur many flexures, different positions, and occafional irritations; it is obvious, that refiftances and ftronger contractions may frequently occur here,

here, to change the direction of the motion ; accordingly we find it frequently changed, and directed from below upwards, in fo far that the contents of the intestines frequently pass into the stomach. But the motions of the intestines are, however, most constantly directed from above downwards, both because they are commonly begun from the ftomach, and becaufe, when inverfions do occur, there is commonly still fo much refistance at the pylorus, and more efpecially at the valve of the colon, as to turn the direction again into its proper course. In the colon, from its polition, structure, and the confistence of its contents, the progress of these is more flow and difficult, and it is therefore here affifted by the longitudinal fibres pecu-

peculiarly difpofed, fo as by their contraction to contribute more to the dilatation of every fucceeding portion of the inteffine.

CCXXVI. The chyle is taken into the lacteals, and moved onwards in thefe in the fame manner (CLXVII.) as the lymph is in the feveral lymphatics in other parts of the body, to which the lacteals are in ftruc-

ture and fituation exactly fimilar.

CCXXVII. The course of the alimentary matters, and the motions by which they are carried on, being now explained; we return to confider the feveral changes which the aliment undergoes in this course.

# Q2 CCXXVIII.



CCXXVIII. In the mouth, if the aliment taken in be of a folid confistence, it is here, as we have faid, subjected to a triture; and if our food is of a foft and moift kind, we are inftinctively directed to take in along with it fome dry matter, as bread, that the whole may be fubjected more certainly to a compleat manducation. By this our aliment is not only more minutely broken down, but is also intimately mixed. with the liquids at the fame time taken in, with the faliva and other fluids of the mouth, and with a quantity of air intangled by these viscid fiuids.

CCXXIX. In this divided and moiftened state, the aliment is taken down into the stomach, where it is farther

farther diffolved, the vegetable matter of it begins to be changed to the nature of animal, and the oily parts of the whole begin to be united with the watery. But these changes by folution, affimilation, and mixture require to be separately confidered.

CCXXX. The folution here, as in other cafes, may be affifted by the mechanical divifion of the folid matter, by the agitation of the diffolving mafs, and by the application of heat, and, with these affiflances, the folution must be performed by the application of a proper menstruum.

CCXXXI. The division of the folid is fometimes affifted by a previous cookery, and commonly by  $Q_3$  the

the manducation we have mentioned; but the human flomach does not feem by any mechanical powers to contribute to this. It gives only a moderate agitation, which, in any cafe, contributes little to mechanical division.

CCXXXII. The degree of heat applied here, being that of the common temperature of the human bo-

dy, may affift the folution; but it is of no confiderable power, and no affiftance is got from any clofenels of the veffel which occurs here. Upon the whole, the affiftances applied here are not confiderable, and the fpeedy folution that takes place muft be chiefly owing to the power of the menftruum.

CCXXXIII.

CCXXXIII. The menftruum that appears here, is a compound of the. liquid matters taken in, of the faliva, and of the gaftric liquors; but in all, or any of these, we do not. readily perceive any confiderable. folvent power; nor, by any artifice,. in employing these out of the body, can we imitate the folutions performed in the stomach.

CCXXXIV. However, from what happens in the ftomachs of certain animals, there is ground to presume, that indeed in every one there is a peculiar folvent. But whether this be a menstruum dividing the folid into integrant parts, and thereby reducing it to a fluid state, or if the solvent here be a peculiar fermentative power, refolving matters



matters more or less into constituent parts, is not clearly perceived.

CCXXXV. The latter is the moft probable, as the circumftances of fermentation very conftantly appear, and as the deviations which at any time appear in the courfe of digeftion appear always to be an excels of fermentation, either acefcent or putrefactive.

CCXXXVI. The bufinels feems to us to proceed in this manner. The fluids of the flomach have the power of fuddenly and powerfully loolening the fixed air of the alimentary matters, which is the firft flep towards putrefaction, and that which most effectually breaks down the texture, and perhaps the mixture

ture of bodies. But we now know, that putrescent bodies are very powerful in exciting an acefcent fermentation in vegetable substances, which the human flomach is hardly ever without; and that this acescency thereforc, in the next place, very constantly fucceeds, and an acid is produced in the ftomach. This acidity makes the effects of the putrefaction disappear; and the acidity in its turn difappears alfo, probably by its being absorbed by or united with the putrefcent and oily matters here prefent; and it is in this man-. ner that we suppose that the animal fluid is produced, and daily renewed by the combination of a fresh portion of acid with putrescent fluids previoufly exifting in the body. The daily production of acid in the human

### igo PITISIOLOGY.

human f heach, and its readily difappearing again, without thewing any morbid effects, renders our doctrine fufficiently probable.

CCXXXVII. This is the affimilation of vegetables that I fuppele to take place, and is begun in the flomach, but is not compleated there ; for we observe that the long retention of the alimentary matters in

the ftomach, whether from the iniolubility of the matter, or from an obftruction of the pylorus, produces a greater degree of acidity, and, ingeneral, the acidity which commonly prevails in the ftomach does not difappear but in the after courfe of the aliment.

#### CCXXXVIII.

CCXXXVIII. It is efpecially the bile, added to the matters which have passed from the flomach into the duodenum, that is fitted to cover the acidity which appeared in the ftomach. It is probable alfo, that the pancreatic and inteftinal liquors contribute to the fame effect; and it is perhaps for the fame purpose, that the lymph is confantly added to the chyle in its course. But, after all, we must reft in the general idea, and own that we do not know exactly how this matter proceeds, nor what the feveral fluids, added to the aliment in the different parts of its course, truly contribute to the changes of it.

CCXXXIX. It is probable however, that, by the mixture mentioned, the

the peculiar fluid which we name the chyle is produced; for, tho' it is certain that a variety of fluid matters may enter the lacteals, and accompany the chyle there, it is ftill probable that there is a peculiar fluid produced by the actions of the ftomach and inteffines, and fuch as becomes the principal ingredient in the -animal fluids afterwards formed that is strictly entitled to that appellation, This chyle does not appear in the ftomach, but first in the duodenum, and more copioufly ftill in .the jejunum and first part of the ileum. It appears indeed in the whole of the ileum, coecum, and colon, but in the last less copiously; all which shews that a particular mixture is neceffary to it, and at the fame time that it is not made at once but fucceffively

ceffively in the course of the inteftines.

CCXL. It remains to fpeak of the mixture of the oily with the watery parts of the aliment. This we cannot well explain; but it is of consequence to observe here, that such a mixture is actually made. It is evident that a large quantity of oil in a separate state is taken in as a part of our aliment, but at the same time no oil commonly appears in a separate state in the mass of blood; it must therefore be united with the other parts of the mass in the way of mixture. Hitherto the physiologifts have hardly mentioned any other means for this union of oil but the application of viscid fluids; but these can occasion only a dif-R fulion,

fusion, and some means of mixture must necessarily be supposed. What thefe however are we do not certainly know. They do not produce their effect in the first passages; for in the chyle, till it enters the fubclavian vein, the oil appears to be only in a diffused state, and probably the perfect mixture is only made in the paffage through the lungs.

CCXLI. It may be proper here to take notice of another matter which conftantly enters into the mixture of animal fluids. This is air, which, by different means, can be extracted in confiderable quantity from every kind of animal matter. What is properly the origin of this, when and where it is infinuated into the animal fluids, and by what means it is either fixed

ed in these or loosened from them, are all questions not yet refolved; but perhaps necessary to be resolved, before we can speak with any confidence of the changes which the animal fluids undergo in different parts of the fystem. We can observe, in the mean time, that a quantity of air is always prefent in the chyle in a very loofe state; that it becomes more fixed in the mass of blood after this has has paffed through the lungs; and that again, in the different fecreted fluids, the air appears to be in fome of them still fixed, and in others much more loofe; and it is probable, that all this has a particular relation to the production and properties of the different fluids of animals...

R 2

CCXLII.

CCXLII. We have now followed the courfe of the aliments, fo far as we can confider them as any ways in a leparate ftate ; but we do not perceive, that, in any part of this courfe, the proper animal fluids are intirely formed : And it is very juftly fuppofed, that the proper mixture or affimilation is not finished till the chyle, mixed with the mass of blood, has undergone

the action of the lungs, through the veffels of which it must almost immediately pass, after entering the fubclavian vein, and seemingly before it is applied to any of the purposes of the animal oeconomy.

CCXLIII. What change the fluids undergo in paffing through the lungs, or by what means the fuppofed

posed changes are produced, after all that has been said, seems still to be very little known.

The mechanical powers of preffure, commonly fpoken of, do not in fact take place, nor are their fuppoled effects any ways confiftent with found philofophy; and, on the other hand, it is very probable; that the changes produced are the effects either of chemi-

cal separatic a or mixture.

What has been fuppofed to be performed in this way by an abforption of air, or of a particular matter from it, is very uncertain in fact, and has led to a ftill more uncertain reafoning.

It is now certain, that a quantity of mephitic air, and perhaps fome  $R_3$  other

other matters, are constantly exhaling from the lungs of living animals, and are carried off by the atmospherical air alternately entering and isfuing from the lungs. This is a pretty certain evidence that fome change of mixture is going on in the fluids paffing through the lungs; but from what particular portion of the fluids the mephitic air proceeds, or what is the effect of its leparation, we know not: And indeed, as we have faid before, what are the effects of the action of the lungs upon the state of the fluids, we are very uncertain. Upon the whole, we still know but little of the production or formation of the animal fluids; and therefore, from the confideration of their formation, we have learned little of their nature; but

but we must now try to difcover what we can of it, by examining these fluids as they are found already formed in the blood-veffels.

#### SECT. IV.

CHAP. II.

#### OF ANIMAL BLOOD.

CCXLIV. The red fluid paffing from the lungs to the left ventricle of the heart, and thence by the aorta and its branches to every part of the body, may be confidered as

a

a mass containing, either formally or materially, every part of the animal fluids, and may therefore be properly called the common mass of blood. This term, however, must be strictly confined to the circulating fluids while they retain their red colour; for, when they lose this, it is always in consequence or some separation of parts. The same red fluid, indeed, as it is found. in the veins; has also fuffered some separation of parts; but as the blood in the veins is never intirely deprived of the whole of any matter that was prefent in the arteries, fo we think the venous blood may still be confidered as a part of the common. mais.

CCXLV. This mass of blood we find

find to be an heterogeneous aggregate; and it will be proper to inquire into the feveral parts of this, before we employ any chemical trials for difcovering the mixture of the whole, or of its parts.

CCXLVI. We difcover the parts of this aggregate chiefly by the fpontaneous feparation of them, which takes place upon their being

drawn out of the veffels of a living animal.

CCXLVII. The feparation commonly proceeds in this manner. Immediately after the blood is drawn out, it exhales a fenfible vapour, and, after fome time, it is found, by that exhalation, to have loft a part of its weight, more or lefs,

lefs, according to the degree of heat it is exposed to, according to the extent of furface by which it is exposed to the air, and probably alfo according to different conditions of the blood itself. The matter thus exhaling may be called the *balitus* or vapour of the blood.

CCXLVIII. Soon after the blood has been drawn out of the veffels,

it lofes its fluidity, and the whole of it concretes into one foft gelatinous mafs; but, after fome time, there ouzes out from this mafs a thin fluid, and as the feparation of this proceeds, the mafs contracts into a finaller bulk, and, in proportion, becomes more denfe.

CCXLIX.

CCXLIX. This is the feparation which almost always takes place, and has, at all times, been observed by physicians. The fluid part is called *ferum*, and the thicker confistent part has been called *cruor*, but more properly the *craffamentum*.

CCL. Both parts feem homogeneous and fimple, but are not. For, if the craffamentum taken from the ferum be laid upon a linen cloth, and water is poured upon it, the water wafhes off a red coloured part, and carries it through the pores of the cloth, and there remains a whitifh, confiftent, but foft and tough mafs, not to be further diminifhed or feparated into parts by any ablution.

A like experiment flews always a like matter prefent in the mass of blood; and, upon feveral occafions, both while the blood remains within the veffels, whether of the living or dead body, and when it is drawn out of the veffels of the living, this matter fpontaneoully separates from the other parts of the blood. It is therefore a part constantly present in the blood. It is what Gaubius, after Malpighi, calls the fibra fanguinis. Mr Senac names it the coagulable lymph, and we shall speak of it under the title of the gluten of the blood. When it appears upon the furface of the blood drawn out of the veffels of living animals, it is called the inflammatory cruft.

CCLI.



CCLI. When the blood is viewed with a microfcope, whether as moving in the veffels of a living animal, or when out of the veffels remaining still fluid, there are certain parts of it which appear of a round figure, and also of a red colour, while the reft is almost colourles. The parts thus diffinguishable by their figure, are called the red globules, and it appears, that the red colour of the whole mass depends upon the presence of these only. It is chiefly thefe parts which are washed off from the crassamentum in the experiment above-mentioned; and we now conclude, that, befides the red globules; the gluten, and a portion of ferum that happens to be entangled in the pores of the con-S creting

· .

creting mass, there is no other matter evident in the crassamentum.

CCLII. The ferum is a transparent fluid of very little colour, and feemingly fimple; but, if it be exposed to a heat of 156 degrees of Fahrenheit's thermometer, it concretes into a firm and almost transparent gelly; and, if this be cut into minute pieces, there exudes from it a thin colourless fluid of a faline taste. In proportion as this fluid is more carefully feparated, the coagulated part becomes more infipid, and in all its properties refembles the gluten feparated from the craffamentum. From hence we are ready to conclude, that the ferum, as obtained by spontaneous separation, consists of a portion of gluten

gluten diffolved in a faline fluid, which we name the SEROSITY.

CCLIII. From the whole that has been faid from (CCXLIV. to CCLII.), it appears, that there are three diffinct portions and kinds of matter in the common mafs of blood, that is, red globules, gluten, and ferofity. What other matters may alfo be there, we fhall confider afterwards; but, in the mean time, fhall fay a little more of each of the parts we have already mentioned.

CCLIV. The red globules have been confidered as an oily matter, and from thence their diffinct and globular appearance has been accounted for; but there is no direct proof of their oily nature, and their ready union with, and diffufibility S 2 in

in water renders it very improbable. Asbeing microscopical objects only, they have been represented by different persons very differently. Some have thought them fpherical bodies, but divisible into fix parts, each of which, in their separate state, were also spherical; but other persons have not observed them to be thus divisible. To many observers, they have appeared as perfectly fpherical, while others judge them to be oblate spheroids or lenticular. To fome they have appeared as annular, and, to others, as containing a hollow veficle. All this, with feveral other circumstances relating to them, very varioully represented, fhew fome uncertainty in microfcopical observations, and it leaves me, who am not conversant in fuch obfervations,

fervations, altogether uncertain with refpect to the precife nature of this part of the blood. The chemical hiftory of it is equally precarious; and therefore, what has been hitherto faid of the production, and charges happening to thefe red globules, we chufe to leave untouched. We shall afterwards fay fomething, with refpect to their general ufe in the animal fystem; and now we shall at-

tempt to explain the caufe of fome changes, which in certain circumftances appear in the colour of the whole mass of blood.

CCLV. We fuppofe that the red globules, when viewed fingly, have very little colour, and that it is only when a certain number of them are laid upon one another, that the colour appears of a bright red; but this alfo has its limits; fo that,  $S_3$  when

when the number of globules laid on one another is confiderable, the colour becomes of a darker red. Upon this fuppolition, the colour of the mass of blood will be brighter or darker as the colouring part is more or less diffused among the other parts of the mass; and we think this appears to be truly the case from every circumstance that attends the changes which have

been at any time obferved in the colour of the blood.

CCLVI. The gluten of the blood, from its refemblance, on the one hand, to the *albumenovi*, and, on the other, to the matter of the folids of animal bodies, we confider as the principal part of animal fluids, as that which is immediately formed of

of the aliment taken in, and as that which is employed in increasing the growth of the folids, or in-repairing their wafte.

CCLVII. But it is well known, that the animal fluids in general, and particularly the gluten, is prone to putrefaction; and that, even in the living body, if fresh aliment be not constantly taken in, and also if

certain excretions which carry off putrefcent matter be not conftantly fupported, a confiderable putrefaction certainly takes place. From hence we are led to think, that fome approach to putrefaction conftantly takes place, even in the moft healthy bodies, and that it appears efpecially in an evolution of faline matter, and that this, taken up by the water

water conftantly prefent, forms the ferofity. We suppose it is this which affords the vapour of the blood; (CCXLVII.) and that it is the ferofity diffolving a portion of the gluten which forms the ferum that appears upon spontaneous separation. (CCXLVII.)

CCLVIII. The faline matters impregnating the ferofity, if we may:

judge from the analyfis of urine, are of various kinds; but; particularly, there is prefent an ammoniacal falt, now well known under the name of the *effential falt of urine*, which, if not originally formed, is at leaft most copiously evolved in animal fluids.

CCLIX. Thefe are our conjecturcs

tures concerning the parts of animal blood; and it remains to fay in what proportion each of them is present in it. This will perhaps be always difficult; and in the mean time we can perceive, that many estimates formerly made could not be exact; as the feveral parts were not properly known; and, while judging chiefly from the appearances upon spontaneous separation, physicians were not aware how much these are affected by the circumftances of extravalation, and by those in which the blood; is placed after being drawn out. There are not yet indeed experiments made to ascertain, with any exactness, the proportion of the feveral parts men-. tioned; but it is probable, that the: red globules make a small. part of the:

the whole; that the gluten, if we confider both what is in the craffamentum and in the ferum, is in much larger proportion, but that the watery portion is the largest of all; and at the same time that this has always a confiderable quantity of faline matter diffolved in it.

CCLX. We would next put the question, By what means the parts of

this heterogeneous mass are kept fo equably diffused among one another, and the fluidity of the whole so conflantly preferved ? This we suppose to be done chiefly by motion and heat, and by the parts disposed to concrete being kept from the contact of any matters to which they might adhere more firmly than they do to the other parts of the blood. The diffused

fused parts we suppose to be present only in those vessels in which a confiderable degree of agitation is conftantly kept up; and we suppose also that the heat always here prefent both diminishes the cohesion of the gluten and increases the folvent power of the ferofity. Experiments made with neutral falts feem to confirm the latter; and it is also probable that the fame folvent power may be increased by a quantity of air that is conftantly intermixed with the mafs of blood while it remains in the veffels, and is under a constant agitation. It is fupposed that an attention to these feveral circumstances will explain most of the cafes of spontaneous separation that occur either in the living or dead body, within the vessels or without them; but the detail would be too long for this place.

CCXLI.

CCLXI. We shall add here a few words on the use of this fingular composition of animal blood which we have been confidering.

It appears evidently, from many circumftances of the animal occonomy, that its functions require a fyftem of veffels conftantly filled and even diftended; but as, at the fame time, these veffels must be open by a multitude of their extremities, if

all the fluids were fuch as could pais by these extremities, the fystem could not be kept filled for a few minutes. It is necessary, therefore, that the fluids should be partly of fuch a fize as that they cannot pais through the smaller vessels, and partly in a diffused state only, which has commonly the same effect. Hence it is, that the red globules under the ordinary

ordinary impetus of the heart, and arteries are strictly confined to certain veffels, and it is probable, that, in the like circumstances, the diffused gluten does not go much farther. This ferves to keep the larger vessels of the fystem constantly filled ; but, on the other hand, the ferofity being fufficiently fluid might be supposed to run off by the many outlets open to it, and thereby to leave the fluids in the larger veffels of a confistence unfit to circulate. This, however, feems also to be obviated by the vifcidity of the groffer parts of the blood, fufficient always to entangle fo much of the more fluid, as may be neceffary to preferve the due fluidity of the whole.

CCLXII.

CCLXII. The heat of the human body, fupported by powers within itfelf, is probably the effect of the motion of the blood, and might have been treated of, when we were confidering that fubject. But, as many perfons fuppofe it to depend, in fome measure, on the nature of the fluids, we have referved it for this place, and here, perhaps, to fay only, that the queftion concerning

the cause of animal heat is not yet folved.

CCLXIII. The opinion of animal heat's being the effect of mixture, is to be little regarded, as the matters fuppofed to be mixed, the place in which the mixture is made, and the other circumftances velating to it, are equally hypothetical,

PHYSIOLOGY. 219 tical, and the whole is ill fupported by any analogy.

CCLXIV. More fpecioufly is animal heat 'fupposed to be the effect of putrefaction, towards which there is certainly fome approach in animal bodies; but the opinion is still very doubtful. For, first, the effect of any degree of putrefaction in producing heat is not well afcertained: Secondly, It is not supported by any analogy, that putrefaction, in the degree to which only it proceeds in living bodies, is capable of producing the heat appearing there: And, lastly, whatever is the degree to which putrefaction proceeds in living bodies, it does not appear that there is any increase of heat correspondent to the increase of putrefaction, and rather the contrary. T 2 CCLXV.

CCLXV. The fuppositions either of mixture, or of putrefaction, as the cause of animal heat, are both of them rejected by this, that the generation of heat in animal bodies is manifeftly dependent on another caufe, that is, the motion of the blood. For the power of generating heat in any animal is not perfect, till the motion of the blood in it is fully established; and, when the generating power is established, we perceive the heat to be increased or diminished, as various causes increase or diminish the motion of blood. In dying animals, the heat grows lefs, as the motion of the blood grows lefs, and when at death this ceases altogether, the heat ceases also, commonly, at least, as soon after death as we can suppose a body of the fame

PHYSIOLOGY. 221 fame bulk to lofe the heat it had acquired.

CCLXVI. This connection between the heat and motion of the blood feems in general to be well proved; and, tho' it may be difficult to reconcile certain appearances to it, we would fo far admit of the fuppofition, as to inquire, in the next place, into the manner in which the motion of the blood may generate heat.

CCLXVII. On this fubject, the moft common opinion is, that the heat is produced by the attrition of the particles of the blood upon on another, or of thefe on the internal furface of the veffels in which they move. But we cannot find any  $\dot{a}$ -T 3 nalogy

nalogy to fupport either the one or the other fuppofition.

The attempt made to fupport the latter fuppolition, by endeavouring to fhew, that upon this the equality of heat in the different parts of the fame body is well explained, deferves little regard, as it is founded on doubtful principles and miftaken facts.

CCLXVIII. The equality of heat in the different parts of the fame body feems to require the generating power to be very generally diffufed over the whole; but it does not feem to require its being precifely equal in every part, as the interpolition of pretty large velfels in every part of the body, and the fpeedy communication of the fluids from

from any one part to every other, will fufficiently account for the e-quality of heat, tho' the generating power fhould be in fome meafureconfined to certain parts only.

However, we take no notice of the fuppofitions which have been made of the generating power's being confined to certain finall portions of the fyftem only. Thefe fuppofitions give no relief in the general theory, and they are not fupported by any particular evidence. The breathing animals are the warmeft; but that they are warmer becaufe they breathe, is not more probable, than that they breathe becaufe they are warmer.

CCLXIX. With refpect to this theory, which deduces animal heat from

from the motion of the blood, we must own, that it is attended with feveral difficulties. It will be difficult to fhow, in for many animals of differentage, fize, and temperament, in which the degree of heat is nearly the fame, that the motion of the blood, in all its circumftances, is alto exactly the fame; or to fhow, in the different animals in which the degree of heat is confiderably different, that the motion of the blood in each is correspondent to the difference of heat. May it not be fupposed, that there is some circumstance in the vital principle of animals which is in common to those of the fame clafs, and of like oeconomy, and which determines the effect of motion upon the vital principle to be the fame, tho' the motion acting

PHYSIOLOGY. 225 ting upon it may be in different

circumstances.

CCLXX. In all we have hitherto faid of animal fluids, we have confidered the common mass of blood as confifting of three parts or three kinds of matter only; but many more have been fupposed to be present in it; and we fhall inquire upon what ground. It is common to fuppose that the aliment or the chyle formed of it is not perfectly affimilated in paffing once only through the lungs; but that, for some time after such paffage, it continues to circulate with the blood under the fame form and of the fame qualities which it had when it first entered the fubclavian, and particularly in this state to furnish the

the milk which is fecreted in the breafts of females. There is however no proper evidence of the chyle's ever appearing in the blood-veffels, and the appearances of it alledged can be otherwife accounted for. The arguments for the fame opinion which are drawn from the confideration of the fecretion of milk are embaraffed with many difficulties.

CCLXXI. It is probable that the animal fluid (CCLV. CCLVI) is in a conftant progrefs, and hardly for a moment flationary, or therefore uniformly the fame over the whole of the common mafs. Some part of it is that which was laft formed, and therefore the neareft to the vegetable matter from which chiefly it was produced; while another part of it is

is that which has remained longeft in the body, and is therefore the neareft to putrefaction. Between thefe two there may be feveral intermediate flates, which however, like the neareft flades of the fame colour, are not diffinguifhable by our fenfes or experiments.

CCLXXII. Befides the difference of matter arifing from the progrefs of the animal fluid, there have been other matters fuppofed prefent in the common mafs, and as commonly conflituent parts of it. Such are a mucous matter, like to the mucous matter of vegetables; and a gelatinous matter, like to that which is extracted by decoction from the folid parts of animals. But there is no evidence of either being formally

ly prefent in the mass of blood, and the supposition is founded on mistaken facts and false reasonings.

CCLXXIII. But it is proper to be observed here, that many extraneous matters may, by different ways, be introduced into the blood-veffels; and that many of the fecreted fluids, iometimes very different from any thing that existed before in the mass of blood, may, by absorption or regurgitation, be again taken into the blood-veffels: But, with regard to all of these, whether extraneous matters, or those produced in the body itself, it is probable that hardly any of them enter into the mixture of the animal fluid, and that they are only diffused in the serofity till they can be

be again thrown out of the bloodveffels by the readieft outlets. The oil of the adipofe membrane is frequently, and perhaps neceffarily reabforbed, and feems to be, befides the lymph, the only reabforbed matter which enters again into the mixture of the animal fluid.

#### SECT. IV.

#### CHAP. III.

### OF SECRETION.

CCLXXV. After thus confidering the parts of the mass contained U in

in the red veffels, we must next confider the feveral fluids which appear in the other parts of the body.

CCLXXVI. All of these we suppose to be derived from the common mass, as they appear in vessels continuous with those of the common mass, and as their appearance ceases when the communication of the vessels containing them, with the fanguiferous vessels, is any how interrupted.

CCLXXVII. The fluids thus derived from the common mafs feem to be produced in confequence of a certain ftructure, with perhaps fome other condition in the extreme veffels through which the fluids pafs; and a part having fuch a ftructure,

18

PHYSIOLOGY. 231 is called a *gland* or *fecretory organ*, the function of which, from the most obvious notion of the manner of it, is called *fecretion*.

CCLXXVIII. The ftructure of the organ, and the manner of its function, feem to me for the most part unknown; at least, what we know or suppose with regard to the ftructure hardly in any case applies

to the explanation of the function.

CCLXXIX. If it any how appeared that the feveral fecreted fluids were all of them previoufly exiftent in the fame forms in the mass of blood, it would not perhaps be difficult to explain what might be ftrictly called a fecretion. But fuch previous existence does not appear;  $U_2$  for,

for, except the matter of exhalation into the feveral cavities of the body, and the matter of urine and of perfpiration, we find no proper evidence of any other fecreted fluids prefent in the mass of blood. We cannot find there, either milk, mucus, or oil, and much less the appearance of many other fluids, which are only found after they have passed thro' certain organs.

CCLXXX. This being the cafe, the confiderations of the phyfiologifts with regard to the velocity of the blood, and other circumftances favouring the feparation of the parts of a fluid which are only diffufed among one another, deferve no attention. The effects of different apertures may go fome length; but we

we can perceive their particular application only in the few cafes of a fimple feparation. In most others, there appears to be a change of mixture ; but we perceive neither the precife changes that are made, nor the cause of them.

CCLXXXI. Till we can difcover thefe more clearly, we may in the mean time obferve, that the action of the veffels of the fecretory organ have a confiderable fhare in determining both the quantity and quality of the fecreted fluid, and that both very often are very little affected by the general flate of the circulation, or by the different conditions of the mafs of blood.

CCLXXXII. It would feem that U 3 no

no other fecretion but those of peripiration and fweat are manifestly increased by the increased action of the heart and arteries, (CLXXXI.) and that most of the other fecretions are increased only by stimulants applied to their organs. These stimulants may be either fuch as are immediately applied externally or internally to the excretory, or perhaps to the fecretory veffels, or they may be fuch as are applied to the fensorium, or to distant parts of the nervous fystem, which by the laws of the animal oeconomy have a connection with the organs of fecretion. These stimulants, at the same time that they act in either of these ways on the fecretory organs, for the most part have no sensible effect on the general state of the circulation of the blood. CCLXXXIII.

·

CCLXXXIII. With respect to the influence of the condition of the common mass of blood upon the several secretions, we prefume that the state of the quantity of the fluids. in general will affect the quantity of every fecretion; but the effects of the quantity of the whole mass are very remarkable only, with respect to the secretions of perspiration, urine, and milk. The qualities of the common mass may also be prefumed to affect the feveral secretions; but the effect of these qualities appear most remarkable in the fàme fecretions of perspiration, urine, and milk, and; even in thefe, the effect seems to depend upon the proportion of water more than upon that of any other matter m



in the common mass. With respect to the other fecretions, we cannot perceive that any of them are increased by a particular matter prefent in the mass of blood, except it be such a matter as stimulates the secretory organ.

CCLXXXIV. The feveral fecretions are frequently obferved to affect each other mutually, fo that

the increase of one diminishes another, and vice versa. This seems to depend either upon a change of determination in the course of the blood (CLXXVIII.) or upon a change in the state of the sluidity of the common mass, or, perhaps, upon a connection established between the different organs of secretion as parts of the nervous system; and, except it

it be in the cafe of perfpiration and urine, we cannot perceive that the effect of the state of one secretion upon that of another depends upon an increase or diminution of any particular matter in the mass of blood.

CCLXXXV. After mentioning these generalities with respect to secretion, we should, perhaps, proceed

in the next place to confider the application of them to the particular fecretions, and alfo to confider more particularly the feveral fecreted fluids; but we omit both thefe fubjects, as we prefume the former will be obvious from what is already faid; and with refpect to the latter, we have not yet a fufficient number of experiments

# 238 PHYSIOLOGY. experiments to proceed any length

in it.

#### SECT. IV.

CHAP. IV.

OF NUTRITION.

CCLXXXVI. Under this title we might confider how the matter both of the fluids and folids of the body is fupplied; but, after what we have formerly faid of the taking in and affimilation of the aliment, we have nothing now to add with refpect to the fupply of the fluids;



fluids; and we therefore confine ourfelves here to confider in what manner the folid parts obtain their increase of matter and growth, or have their occasional waste repaired.

CCLXXXVII. There is no doubt that the folids are formed of the fluid prepared from the aliment in the manner we have faid; but it is required now to fay what portion of the fluids is employed in nourifhing the folids, by what channels the nourifhment is conveyed to them, and, being applied there, how from fluid it becomes folid.

CCLXXXVIII. With regard to the first question, we have no doubt in afferting, that in oviparous animals, it

•

it is the *albumen ovi* that is employed in nourifhing the chick; and we prefume that it is an analogous fluid which is employed in nourifhing the bird during the whole time of its growth. We think the analogy may be fafely applied with refpect to all animals, the folid matter of which is of the fame kind with that of the oviparous.

CCLXXXIX. This analogous fluid we take to be the gluten of the blood, properly diluted and freed from any adhering faline matter.

CCXC. To determine in what manner this nutritious fluid is applied to the nourifhment of the fc'i's, it is neceffary to confider what are the fimple fundamental folids

PHYSIOLOGY. 241 lids, of which all the others are formed.

CCXCI. It feems to be the opinion of the greater part of modern anatomists, that the folid parts confift entirely of a cellular texture, of various denfity in the different parts; and indeed, the structure of the greatest part of the folids is evidently of this kind. But, at the fame time, it is also true, that a fibrous ftructure is to be observed almost every where in the body. It appears in the medullary fubftance of the brain and nerves, in the muscles and tendons, in the arteries, in the excretories of the glands, in the lymphatic vessels, in the alimentary canal, in the uterus and bladder of . urine, in the ligaments, in most mem-X branes,

branes, and it is to be feen in those membranes which are afterwards changed into bones, especially whilst this change is going on.

CCXCII. From this view of the univerfality of a fibrous ftructure in animal bodies, we are difpofed to believe, that thefe fibres are the fundamental part of animal folids, that they are the primordial ftaminal part of animal bodies, and that the cellular texture is, for the most part, an accretion formed upon these fibres.

The confideration of the ftructure and growth of vegetables feems to illuftrate and confirm this opinion.

CCXCIII. At the fame time, from the fibrous parts (CCXCI.) being evidently,

vidently, in most instances, parts of the nervous fystem, and from the gradual formation of the foetus, in which the nervous fystem is first formed, we think it probable, that the whole of the fibres in the different parts of the body, are a continuation of the nerves; and this again will lead to the conclusion, that the nourisimment of the fost and homogeneous folid every where is

conveyed to it by the nerves.

CCXCIV. This fuppofes alfo, what is otherwife probable, that the cortical part of the brain, or common origin of the nerves, is a fecretory organ, in which the gluten of the blood, being freed from all faline matter before adhering to it, becomes fit for the nourifhment of the X = 2 folids,

folids, and being poured in a fufficiently diluted flate upon the origin of the nerves, it is filtrated along the fibres of these, and is thus conveyed to every flaminal fibre of the fystem. We suppose, at the fame time, that the medullary, or what may be called the folid matter of the nerves, is in the living body conftantly accompanied with a fubtile elastic fluid, which fits them for being the organs of fense and motion, and which probably is alfo the means by which the nutritious fluid is carried on in the fubflance of the nerves, from their origin to their extremities.

In what manner the nutritious fluid, thus carried to the feveral parts, is there applied, fo as to increafe the length of the nervous fibre

fibre itfelf, or to form a cellular texture upon its furface; and in what manner from fluid it becomes folid, we cannot explain; nor can thefe particulars be explained upon any other fuppofition that has been formed with refpect to nutrition.

.

CCXCV. It is probable that, for a certain time, at its firft beginning, the growth of animal bodies proceeds in the fame manner as that of vegetables; but it is evident, that, at a certain period, in the growth of animals, a different oeconomy takes place, and that, afterwards, the growth feems to depend upon an extension of the arteries in length and wideness, by the blood propelled into them by the powers (CLVI. CLIX.) It may be fup-X 3 posed,

posed, that this extension of the arteries is applied to every fibre of the body, and that, by the extension of thefe, it gives an opportunity to the application and accretion of nutritious matter; to the growth therefore of the fibre itself, and to the growth of cellular texture on its furface. Perhaps the fame extension of the arterial fystem gives occasion to the fecretion of fluids, which, poured into the cellular texture already formed, according to the disposition of these fluids to concrete more or less firmly, gives the different degrees of denfity and hardness which appears in different parts of the body.

CCXCVI. By this extension of the arterial system, the several parts of the body are gradually evolved, fome

# PHYSIOLOGY. 247.

fome of them fooner, others later, as by the conflitution of the original stamina, or after occurrences, they are feverally put into the conditions (CLXXVII. CLXXVIII.) by which they are more or lefs exposed to the impetus of the blood, and fitted to receive a greater quantity of it. But as the parts by these caufes first evolved will increase the most in the density of their folid parts, they will therefore more and more refift their further growth, and by the fame refiftance, will determine the blood with more force, and in greater quantity, into the parts not then fo far evolved. Hence the whole fyftem will be at length evolved, and every part of the folids will, in respect of density and resittance, be in balance with every

every other, and with the forces to which they are feverally exposed.

(CCXCVII.) The extension of the arteries (CCXCV.) depends upon the refiftances which occur to the free transmission of the blood thro' them, as in (CLXX.), and further; from a refistance in the veins. For, as a confiderable portion of the blood, by (CCLXI.), does not commonly pais into the finaller branches of the arteries, but must pass very entirely into the veins, fo thefe, by their capacity conftantly diminishing, as they approach nearer to the heart, and by their coats being of a denfity and firmness sufficient to prevent further dilatation, confiderably refift the free paffage of

PHYSIOLOGY. 249 of the blood from the arteries intothem.

CCXCVIII. While thefe refiftances continue, the arteries, and with them almost every fibre of the body, must be extended at every fystole of the heart, and with this extension, the growth of every part will proceed; but, as every part, by its receiving an addition of folid matter, becomes more denfe and rigid, fo it is less eafily extended, and perhaps less readily receives an accretion of new matter than before. Hence it is, that the more the body grows, it admits of any additional growth more flowly; and, unlefs the extending powers increase in the fame proportion with the increasing denfity of the folids, there must be

Э.

a period at which these two powers. will balance each other, and the growth will proceed no farther. But, as it is evident, that the bulk and weight of the heart, and probably therefore, its force, does not increase with the increasing bulk of the body, and that the action of the heart is the principal extending power in the system; it is also plain, that the extending power does not increase in the same proportion with the increasing density of the folids; and therefore, that thefe two powers will, at a certain period, come to balance each other.

CCXCIX. But, not only is the force of the heart thus conftantly diminishing, with respect to the refistance of the arteries, but the force. of

of the heart, though it were still fubfifting, has, from other caufes, lefs effect in extending the arteries. The blood is more confined to the arteries, and extends them further in proportion to the refiftance in the veins, as in (CCXCVII.); and this refiftance in the veins, and the extension of the arteries depending upon it, will be more or lefs, according to the respective density of these two sets of vessels. But it appears from the experiments of Sir Clifton Wintringham, that the denfity and firmness of the veins, with respect to their correspondent arteries, is much greater in young animals than in old; and thence it appears, that, during the growth of animals, the ar-

teries are acquiring an increase of denfity in a greater proportion than the

the veins are at the fame time; and therefore, that the refiftance in the veins with refpect to the arteries, must be constantly diminishing; that the veins will therefore receive a a greater proportion of blood; that in the fame proportion the arteries will be less extended; and, *laftly*, that the diminished refistance in the veins, concurring with the diminished force of the heart, will the

fooner bring the increasing rigidity of the arteries, and therefore, of every fibre of the body, to be in balance with the extending powers; at least, fo far as to prevent their producing any further growth.

CCC. This account of the change of the refiftances in the arteries and yeins

veins, with refpect to one another, is agreeable to phaenomena, which fhew that the arteries are larger, and contain more blood in proportion to the veins in young animals, than in old; that arterial haemorrhagies occur most frequently in young perfons, and that congestions in the veins, with haemorrhagies, or hydropic effusions depending upon

them, occur most frequently in old age.

CCCI. It is probable, that the refiftance both of arteries and veins, goes on increafing, while the force of the heart is not increafing at the fame time; but it appears alfo, that, from the diminished force of the heart, and the compression which the finaller vessels are constantly ex-

pofed

posed to from the distension of the larger, the action of the muscles, and other causes; the number of fmall veffels, and therefore the capacity of the whole fystem, is conflantly diminishing fo much, that the heart may still, for some time, be fufficient for the circulation of the blood. But, while the refiftances in the veffels are constantly increasing, the irritability of the moving fibres, and the energy of the brain, are at the fame time constantly diminishing; and therefore the power of the heart must at length become unequal to its tafk, the circulation must cease, and death enfue.

CCCII. The unavoidable death of old perfons is thus, in part, accounted

ed for; but it is, however, still probable, that the fame event proceeds chiefly from the decay and total. extinction of the excitement or vital power CXXXVI. of the nervous fystem, and that from causes very much independent of the circulation of the blood, and arifing in the nervous fystem itself, in confequence of the progress of life. This feems to be proved by the decay of fense, memory, intellect, and irritability, which constantly takes place, as life advances beyond a certain period.